

Progress in Classical and Quantum Variational Principles.*

C.G.Gray [†] G.Karl [‡]

Physics Department, University of Guelph,
Guelph, Ontario, Canada, N1G 2W1

V.A.Novikov [§]
ITEP, Moscow, Russia

*Reports on Progress in Physics (2004)

[†]cgg@physics.uoguelph.ca

[‡]gk@physics.uoguelph.ca

[§]novikov@heron.itep.ru

“...the most beautiful and important
discovery of Mechanics.”

Lagrange to Maupertuis (November 1756)

Abstract

We review the development and practical uses of a generalized Maupertuis least action principle in classical mechanics, in which the action is varied under the constraint of fixed mean energy for the trial trajectory. The original Maupertuis (Euler-Lagrange) principle constrains the energy at every point along the trajectory. The generalized Maupertuis principle is equivalent to Hamilton's principle. Reciprocal principles are also derived for both the generalized Maupertuis and the Hamilton principles. The Reciprocal Maupertuis Principle is the classical limit of Schrödinger's variational principle of wave mechanics, and is also very useful to solve practical problems in both classical and semiclassical mechanics, in complete analogy with the quantum Rayleigh-Ritz method. Classical, semiclassical and quantum variational calculations are carried out for a number of systems, and the results are compared. Pedagogical as well as research problems are used as examples, which include nonconservative as well as relativistic systems.

Contents

1	Introduction and History	4
2	Action Principles of Classical Mechanics	6
2.1	Statements of Action Principles	6
2.2	Derivation of Action Principles	8
2.3	Semiclassical Mechanics from Action Principles	11
3	Practical Use of Variational Principles. Pedagogical Examples	11
3.1	The Quartic Oscillator	12
3.2	The Spherical Pendulum. Precession of Elliptical Orbits . . .	14
4	More Complicated Examples. Research Problems	17
4.1	The x^2y^2 oscillator	18
4.1.1	Perturbative Shift of Variational Results	22
4.1.2	Splitting of Degenerate States	24
4.2	The “Linear Baryon”	26
4.3	He Atom Inside C_{70} Cage	29
5	Nonconservative Systems	34
5.1	Action Principles	34
5.2	Adiabatic Invariants and the Hannay Angle	36
5.2.1	Bead Sliding on Slowly Rotating Horizontal Ring . . .	39
5.2.2	The Foucault Pendulum	41
5.2.3	Nonintegrable Systems	44
5.3	The Classical Hellmann - Feynman Theorem	46
6	Variational Principles in Relativistic Mechanics	48
7	Classical Limit of Quantum Variational Principles	53
8	Summary and Conclusions	58
9	Acknowledgements	60
10	Appendix I. Variational Notation and Reciprocal Principles	60
11	Appendix II. Equivalence of the Maupertuis and General Maupertuis Principles	60

1 Introduction and History

Variational Principles have a long and distinguished history in Physics. Apart from global formulations of physical principles, equivalent to local differential equations, they also are useful to approximate problems too difficult for analytic solutions. In the past century quantum variational calculations have been ubiquitous as an approximate method for the ground state of many difficult systems [1]. In recent years there has been some progress in using classical variational principles (Action Principles) to approximate the motion of classical systems such as classical molecules. We review recent developments in this area of Classical Mechanics, although we also overlap with Quantum Mechanics. In particular we shall also discuss the use of the quantum variational principle for excited states, and the connection to classical action principles.

In Classical Mechanics, variational principles are often called Least Action Principles, because the quantity subject to variations is traditionally the Action. To confuse matters there are two classical Actions, corresponding to two main Action Principles, which are called respectively, Hamilton's Action (S) and Maupertuis' Action (W); these differ from each other (they are related by a Legendre transformation), and the notation S, W is not universal (we follow current usage but in fact some authors use the same symbols reversed [2]). Both Actions, S and W , have the same dimensions (i.e. energy $\times$ time, or angular momentum). Maupertuis' Least Action Principle is the older of the two (1744) by about a century, and as we shall discuss, it is inconveniently formulated in all textbooks [3]. For clear statements of these old action principles, see the textbooks by Arnold [4], Goldstein et al [5], and Sommerfeld [6]. We shall review these Action Principles in the next section. Some mathematics and notation required to deal with variational problems are summarized in Appendix I. The distinguishing feature of variational problems is their global character [7]. One is searching for the function which gives a minimum (or stationary value) to an integral, as opposed to local extrema, where one is searching for the value of the variable which minimizes (or makes stationary) a function. Of course this dichotomy is not so sharp in practice, as the global problem is equivalent to a differential equation (the Euler-Lagrange equation) which is of course local. The relation has even found its way into modern parlance [8]: "Think globally, act locally!"

In Quantum Mechanics, where (usually) the aim is for approximations to

the energy by choosing a wave-function, one speaks of trial wave-functions, which are optimized according to a Variational Principle. In Classical Action Principles, the equivalent notion is that of a virtual path or trajectory, which should be optimized in accordance with the Action Principle. For uniformity we shall talk of Trial Trajectories in the classical domain, to emphasize the similarity of these ideas.

As noted above, Maupertuis' Action Principle was formulated in precise form first by Euler and Lagrange some two and a half centuries ago; for the history, see the books by Yourgrau and Mandelstam and by Terral [9]. However, even the formulation of these eminent mathematicians remained problematic, as emphasized by Jacobi, who stated in his Lectures on Dynamics: “In almost all text-books, even the best, this Principle is presented so that it is impossible to understand” [10, 11]. To understand at least in part the problem (see [12] for a detailed discussion), it is sufficient to note that Maupertuis' Principle assumes conservation of energy, whereas a well formulated principle, like Hamilton's Principle, implies energy conservation. As we shall see, it is possible to reformulate Maupertuis' Principle in a more general form to remedy this problem and also make it more useful in applications [12, 13]. It should be noted that Hamilton's Principle does not suffer from any such drawbacks and is useful both conceptually (to derive equations of motion) and as a tool for approximations. The reformulated Maupertuis Principle has the advantage of being closely related to the classical limit of Schrödinger's Variational Principle of wave mechanics, and thereby lends itself easily to semiclassical applications.

Even though we emphasize two Action Principles for Classical Mechanics, it is worth noting that there are many other formulations and reformulations of these principles. For a list of references on this point see ref.[12]. We note in particular the elegant work of Percival in the 1970's on variational principles for invariant tori (see ref.[14] for a review). The emphasis on Maupertuis and Hamilton reflects the two main approaches of mechanics, one based on the Lagrangian L and the other based on the Hamiltonian H .

In the above and in section 2 we use the traditional terminology for the variational principles, i.e. “least” action principles, but, since Jacobi's work it has been recognized [4, 11] that the action is in fact stationary in general for the true trajectories. This means the first-order variation vanishes, and the action may be a minimum, a maximum or a saddle, depending on the second-order variation [15]. Hence “stationary” action principles would be more accurate terminology. Similar remarks apply to the general Maupertuis

and reciprocal variational principles discussed in section 2.

In sections 2-4 we restrict ourselves to conservative holonomic systems. Various problems are solved using classical and semiclassical variational methods and some comparisons are made with the results from the quantum variational method for excited states [16]. In section 5 we discuss nonconservative (but nondissipative) systems, and in section 6 we mention briefly particular nonholonomic systems (which have velocity constraints), in connection with relativistic systems. The relations between classical and quantum variational principles (VP's) are discussed in section 7.

2 Action Principles of Classical Mechanics

2.1 Statements of Action Principles

Hamilton's Least Action Principle (HP) states that, for a true trajectory of a system, Hamilton's Action S is stationary for trajectories which run from the fixed initial space-time point $A \equiv (q_A, t_A)$ to the fixed final space-time point $B \equiv (q_B, t_B)$,

$$(\delta S)_T = 0, \quad (2.1)$$

where $T = t_B - t_A$ is the duration. Here the (Hamilton) action S is the time integral of the Lagrangian $L(q, \dot{q})$ from the initial point to the final point on the trial trajectory $q(t)$,

$$S = \int_{t_A}^{t_B} dt L(q(t), \dot{q}(t)),$$

where $q(t)$ is the generalized coordinate, $\dot{q}(t) \equiv dq/dt$ the generalized velocity and t the time. In practice we choose $t_A = 0$ and $t_B = T$ for convenience. In general q stands for the complete set of independent generalized coordinates $q_1, q_2, \dots, q_f$, where f is the number of degrees of freedom [17]. In (2.1) the constraint of fixed T is indicated explicitly, but the constraint of fixed end-positions q_A and q_B is left implicit. The latter convention for the end-positions is also followed below in (2.2), (2.3), etc. From the Action Principle (2.1) one can derive Lagrange's equation(s) for the trajectory, which we shall not do in this review; see for example [4, 5, 6].

The reformulated (or General) Maupertuis Least Action Principle (GMP) states that, for a true trajectory Maupertuis' Action W is stationary on trajectories with fixed end-positions q_A and q_B and fixed mean energy $\bar{E}$:

$$(\delta W)_{\bar{E}} = 0. \quad (2.2)$$

Here Maupertuis' Action W is given by

$$W = \int_{q_A}^{q_B} p dq = \int_{t_A}^{t_B} 2K dt,$$

where $p = \partial L / \partial \dot{q}$ is the canonical momentum, and in general $p dq$ stands for $p_1 dq_1 + \dots + p_f dq_f$. The second form for W is valid for normal systems, for which $L = K - V$ and K is quadratic in the $\dot{q}$'s, where K is the kinetic energy and V the potential energy. The mean energy $\bar{E}$ is the time average of the Hamiltonian $H(q, p)$ over the trial trajectory of duration T ,

$$\bar{E} = \frac{1}{T} \int_{t_A}^{t_B} dt H(q(t), p(t)),$$

from the initial position q_A to the final position q_B which are common to all trial trajectories. Note that $\bar{E}$ is fixed in (2.2) but T is not, the reverse of the situation in (2.1). We use the notation $\bar{E}$ rather than $\bar{H}$ because in the early sections we restrict [18] ourselves to dynamical variables where the Hamiltonian is equal to the energy $K + V$; the general case is discussed in note [19] and later sections.

The Reciprocal Maupertuis Principle (RMP), or principle of least (stationary) mean energy, is

$$(\delta \bar{E})_W = 0. \quad (2.3)$$

For stationary (i.e. steady-state or bound) motions (periodic, quasiperiodic, chaotic) (2.3) is the classical limit [12] of the Schrödinger Variational Principle of Quantum Mechanics: $(\delta \langle H \rangle)_n = 0$, (see also sec.7). The connection is clear intuitively: for bound motions, in the large quantum number limit the expectation value of the Hamiltonian becomes the classical mean energy and the quantum number n becomes proportional to the classical action W over the motion. This connection enables the RMP to lend itself naturally

to semiclassical applications (sec. 2.3). For periodic and quasiperiodic motions the Reciprocal Maupertuis Principle (2.3) is equivalent [12] to Percival's principle for invariant tori [14, 20]. The RMP is, however, more general and is valid for chaotic motions, scattering orbits, arbitrary segments of trajectories, etc. Percival's principle has been derived by Klein and co-workers from matrix mechanics [21] in the classical limit. Hamilton's Principle also has a reciprocal version [12] - see next section. For a summary of reciprocity in variational calculus, see Appendix I.

The textbook Euler-Lagrange version of Maupertuis' Principle differs from (2.2); the constraint of fixed mean energy $\bar{E}$ is replaced by one of fixed energy E ,

$$(\delta W)_E = 0. \quad (2.4)$$

This is not erroneous, since a true trajectory does indeed conserve energy, but it is the source of the inconveniences referred to by Jacobi [11]. Because, as we show in the next section, the GMP is equivalent to the HP, and therefore to the equations of motion from which energy conservation follows, it will become clear that energy conservation is a consequence of the GMP (2.2), rather than an assumption as in the original MP (2.4).

It is well known that conservation of energy is a consequence of symmetry under time translation (Noether's theorem), either via the Lagrangian and equations of motion [22], or from the action and Hamilton's Principle [23]. Similarly energy conservation can be derived directly from the GMP [12].

In the early sections we assume $L = K - V$ and that V is time-independent (conservative). Later these restrictions are relaxed (see sections 5 and 6).

2.2 Derivation of Action Principles

Within Mechanics one can show that different Action Principles such as (2.1), (2.2) and (2.3) are equivalent to each other. This we sketch below (see [12] for more details). Alternatively, one can show [4, 5, 6] from Hamilton's Principle (2.1) that the Lagrange equations of motion, in some coordinates are essentially Newton's equations in the same set of coordinates. Reference [9] shows that (2.4) is equivalent to Newton's equations, quoting an argument due to Lagrange, and in [12] we derive (2.3) from similar arguments. On a more lofty level one can derive the Action Principles from Quantum Mechanics in the classical limit. Dirac and Feynman [24] have derived Hamilton's

Principle from a path integral formulation of Quantum Mechanics, and similarly the RMP (2.3) can be derived from wave mechanics (see [12]). Further connections between quantum and classical variational principles are given in sec. 7.

The equivalence of Hamilton's Principle (2.1) with the General Maupertuis Principle (2.2) is worth spelling out in a little more detail. (The equivalence of the MP with the GMP is discussed in Appendix II). The starting point of the derivation is the Legendre transformation between the Lagrangian of the system and the corresponding Hamiltonian,

$$H(q, p) = \sum_n p_n \dot{q}_n - L(q, \dot{q}). \quad (2.5)$$

Here $n = 1, 2, \dots, f$ runs over the various degrees of freedom, and q stands for $(q_1, \dots, q_f)$ and p for $(p_1, \dots, p_f)$. Upon integration over time along a trial trajectory starting at $A \equiv (q_A, 0)$ and ending at $B \equiv (q_B, T)$ (where we could have $q_B = q_A$ for a closed trajectory), and with the definitions given above we obtain

$$\bar{E}T = W - S, \quad (2.6)$$

where T is the duration along the trial trajectory. Thus W and S are also related by a Legendre transformation. If we now vary the trial trajectory and the duration T , keeping q_A and q_B fixed, and compute the variation of the different terms in (2.6), we obtain

$$\delta S + \bar{E}\delta T = \delta W - T\delta\bar{E}, \quad (2.7)$$

which is the relation between the different variations. For conservative systems, near a true trajectory Hamilton's Principle is equivalent to the vanishing of the left hand side. In fact a vanishing left hand side is the unconstrained form [25] of Hamilton's Principle (the UHP - see [12] for proof):

$$\delta S + E\delta T = 0, \quad (2.8)$$

where we have used the fact $\bar{E} = E$ on a true trajectory for conservative systems. Eq (2.8) has the unconstrained form $\delta(S + \lambda T) = 0$, with $\lambda = E$ (the energy of the true trajectory) a constant Lagrange multiplier. For fixed T (i.e. $\delta T = 0$), we recover the HP $(\delta S)_T = 0$, and for fixed S (i.e. $\delta S = 0$) we obtain the Reciprocal Hamilton Principle (RHP)

$$(\delta T)_S = 0. \quad (2.9)$$

The RHP (principle of least time) is discussed in detail in ref.[12].

The right hand side of (2.7) (which therefore must also vanish near a true trajectory) gives the unconstrained form of Maupertuis' Principle (UMP) for conservative systems:

$$\delta W - T\delta\bar{E} = 0. \quad (2.10)$$

This has the unconstrained form $\delta(W + \lambda\bar{E}) = 0$, with $\lambda = -T$ (the duration of the true trajectory) a constant Lagrange multiplier. For fixed $\bar{E}$ (2.10) gives the GMP of equation (2.2) while for fixed W it gives the reciprocal, the RMP of equation (2.3). An alternative derivation of the UMP (2.10) is given in ref.[12] with traditional variational procedures, employing the general first variation theorem of variational calculus, which includes end-point variations [15, 29].

Equation (2.7) connecting the different variations is reminiscent of equations of thermodynamics, where two conditions are derived from a single equation. There are many other analogies with thermodynamics, e.g. two actions in mechanics analogous to two free energies in thermodynamics, with a Legendre transform relation in each case [26], adiabatic processes and reversible processes in both mechanics and thermodynamics, and reciprocal variational principles in both mechanics and thermodynamics (see Appendix I). Beginning with Helmholtz [9], many of these analogies have been explored by various authors [27].

As we have stressed, we hold the end-positions q_A and q_B fixed in all the variational principles discussed above. It is possible to relax these constraints by generalizing further the UHP and UMP (see notes [25], [26], and [124]), but we shall not need these generalizations for the applications we discuss.

The derivation we have sketched shows that the four principles (2.1), (2.2), (2.3), (2.9) are equivalent to each other. Mathematically, the Hamilton Principles can be regarded as Legendre transformations of the Maupertuis Principles, since the Legendre transformation relation (2.6) allows us to change independent variables from $\bar{E}$ to T . Thus in Classical Mechanics we have a set of four variational principles that are symmetric under Legendre and reciprocal transformations:

$$(\delta\bar{E})_W = 0 \xleftrightarrow{\text{reciprocity}} (\delta W)_{\bar{E}} = 0 \xleftrightarrow{\text{Legendre}} (\delta S)_T = 0 \xleftrightarrow{\text{reciprocity}} (\delta T)_S = 0.$$

After a short interlude on action principles in semiclassical mechanics we proceed to discuss some simple examples.

2.3 Semiclassical Mechanics from Action Principles

The RMP (2.3) is the basis of a very simple semiclassical quantization method. As we shall see, the solution of (2.3) yields, for the true trajectory or approximations to it, $\bar{E}$ as a function of the action W , $\bar{E}(W)$. For bound motions we then assume the standard Einstein-Brillouin-Keller (EBK) [28] quantization rule for the action over a cycle

$$W = (n + \alpha)h, \quad (2.11)$$

where $n = 0, 1, 2, \dots$, α is essentially the Morse-Maslov index (e.g. $\alpha = 1/2$ for a harmonic oscillator), and $h \equiv 2\pi\hbar$ is Planck's constant. (EBK (or torus) quantization is the generalization of Bohr-Sommerfeld-Wilson quantization from separable to arbitrary integrable systems). Thus we have expressed $\bar{E}$ as a function of the quantum number n , $\bar{E} = \bar{E}_n$. For multidimensional systems, (2.11) is applied to each action W_i in $\bar{E}(W_1, W_2, \dots)$, so that we get $\bar{E}_{n_1, n_2, \dots}$ as a function of the quantum numbers n_i . This method has been found to be reasonably accurate even for nonintegrable systems, where strictly speaking the good actions W_i do not exist. The UMP (2.10) can also be used to obtain semiclassical expressions for $\bar{E}_{n_1, n_2, \dots}$. Examples are given later in sections 4.1 - 4.3.

3 Practical Use of Variational Principles. Pedagogical Examples

Here we discuss how to use variational principles to solve practical problems. We use the so-called direct method [29] of the calculus of variations (e.g. Rayleigh-Ritz), which operates directly with the variational principle and makes no use of the associated Euler-Lagrange differential equation. Historically [30], the method originates with Euler, Hamilton, Rayleigh, Ritz and others (see [12, 29] for some early references). We assume a trial solution, which contains one or more adjustable parameters a_i . We then use the variational principle to optimize the choice of the a_i . Generally speaking,

the more parameters a_i , the better the solution. The direct method is well known in quantum mechanics, but is also very useful in classical mechanics, as we shall see. We give a few simple examples in this section, mainly for bound motions; further simple examples, including for scattering problems, are given in refs. [12, 13] and in the references therein .

The simplest example is the free motion of a particle of unit mass on the line segment $x = 0$ to $x = 1$. With the traditional Maupertuis principle (2.4) once we fix the energy $E = v^2/2$ there is essentially nothing left to vary. With the GMP (2.2) or its reciprocal (2.3) we can choose a variety of trial trajectories. For example we can take velocity v_1 on the sub-segment $0 < x < 1/2$ and velocity v_2 on the sub-segment $1/2 < x < 1$. The action W is $(v_1 + v_2)/2$ and the mean energy $\bar{E}$ is $v_1 v_2/2$, where we can vary v_1 to change W while keeping the mean energy fixed. We then find that $v_1 = v_2$, a simple example of energy conservation for this particular trial trajectory. By iteration we can now see that energy is conserved for the whole trajectory on the segment $(0, 1)$, as a consequence of the General Maupertuis Principle (2.2).

3.1 The Quartic Oscillator

An instructive example is the one-dimensional (1D) quartic oscillator, with Hamiltonian

$$H = \frac{p^2}{2m} + \frac{1}{4}Cx^4. \quad (3.1)$$

Using a harmonic oscillator trial trajectory with

$$x(t) = A \sin \omega t, \quad p(t) = mA\omega \cos \omega t \quad (3.2)$$

we find for a complete cycle with period $T = 2\pi/\omega$

$$\bar{E} = \frac{\omega}{4\pi}W + C \frac{3W^2}{32\pi^2 m^2 \omega^2}, \quad (3.3)$$

$$W = \pi\omega mA^2.$$

From (3.3), with the use of the RMP (2.3) we can obtain the “best” frequency ω_0 solving $(\partial \bar{E}/\partial \omega)_W = 0$, which can be substituted into (3.3) to obtain $\bar{E}$

$$\omega_0 = \left[\frac{3CW}{4\pi m^2} \right]^{1/3}, \quad \bar{E} = \frac{1}{2}(C/m^2)^{1/3}(3W/4\pi)^{4/3} \quad (3.4)$$

and the period T as a function of mean energy:

$$T(\bar{E}) = 2\pi[m^2/(2C\bar{E})]^{1/4}. \quad (3.5)$$

This variational estimate for the period is in error by 0.75% when compared with the exact result, as noted in reference [13]. Systematic improvements can be obtained by including terms $B \sin 3\omega t$, $C \sin 5\omega t$, etc., in the trial trajectory $x(t)$; see [13] for a related example.

We have used the RMP $(\delta\bar{E})_W = 0$ here since (3.3) made it very convenient to do so. The GMP $(\delta W)_{\bar{E}} = 0$ and the HP are also viable for this problem, but the original MP $(\delta W)_E = 0$ is not, since the constraint of fixed E leaves essentially no freedom for variation for 1D problems [31]. One can get around this by relaxing the constraint via a Lagrange multiplier, but this is then equivalent to the GMP (see App. II) or the UMP. In 2D etc., the constraint of fixed E does allow some freedom for variation, but the constraint is very cumbersome and the GMP is always much more convenient.

The use of the action principle (2.3) to obtain the mean energy as a function of the action (3.4) or the period as a function of mean energy (3.5), with harmonic oscillator trial trajectories, is very similar to the use of the variational principle in quantum mechanics to estimate the energies of states of the quartic oscillator with harmonic oscillator trial wave-functions. We show this explicitly for the eigenstates of the Hamiltonian (3.1). In both cases an optimum frequency for the trial harmonic case is sought and the final errors are also similar. We start by computing the quantum expectation value of (3.1) using a harmonic oscillator state with trial frequency ω ,

$$\langle n|H|n \rangle = \frac{1}{2}(n + \frac{1}{2})\hbar\omega + \frac{3C}{8}(\frac{\hbar}{m\omega})^2(n^2 + n + \frac{1}{2}), \quad (3.6)$$

and find the optimum frequency ω_0 by demanding the vanishing of the derivative of $\langle H \rangle$ with respect to the frequency. We substitute this frequency in (3.6) to estimate the energy of excited states

$$\bar{E}_n = \frac{3}{4}(\hbar)^{4/3}(\frac{3C}{2m^2})^{1/3}(n^2 + n + \frac{1}{2})^{1/3}(n + \frac{1}{2})^{2/3}. \quad (3.7)$$

Taking the asymptotic limit of (3.7) for $n \gg 1$ and replacing $2\pi\hbar n$ by W we obtain the second equation (3.4). This is not as trivial as it seems, since (3.7) was obtained from the quantum variational principle (for an arbitrary state [16]), while (3.4) was obtained from a classical action principle.

This example illustrates neatly that the two variational principles are related: the Reciprocal Maupertuis Principle is the classical limit of the Schrödinger Variational Principle (see sec.7 for further discussion). The example illustrates also that the classical action principle can be used as a method of approximation, and semiclassically to estimate E_n as described in sec.2.3. According to (2.11), we quantize $\bar{E}(W)$ in (3.4) semiclassically by the replacement $W \rightarrow (n + \frac{1}{2})h$; as just discussed, this leads to agreement asymptotically ($n \rightarrow \infty$) with the quantum result (3.7). Reference [16] discusses the similar relation between quantum and classical variations for a different one-dimensional system, with a linear potential. The application of classical action principles to other simple systems is discussed in reference [13]. The application of quantum variations for excited states has a long history: the earliest reference is McWeeny and Coulson, who applied it to the quartic oscillator described above [32] more than fifty years ago (for later references see [16]).

3.2 The Spherical Pendulum. Precession of Elliptical Orbits

For a pendulum with two degrees of freedom (θ, ϕ) , there is an additional term in the kinetic energy compared to that of a plane pendulum, i.e.

$$K = \frac{1}{2}mL^2(\dot{\theta}^2 + \sin^2\theta\dot{\phi}^2) , \quad V = mgL(1 - \cos\theta) , \quad (3.8)$$

where θ is the polar angle (measured from the downward vertical axis) and ϕ is the azimuthal angle, m is the bob mass, L the length, and g the gravitational acceleration.

Using the axial symmetry around the z (vertical) axis, we introduce the cylindrical coordinate $\rho = (x^2 + y^2)^{1/2} = L \sin\theta$, and the angular momentum component $l_z = m\rho^2\dot{\phi}$, which is a constant of the motion. We also expand $\cos\theta$, keeping up to quartic terms, which are $O(L^{-2})$. This gives

$$K = \frac{1}{2}m\dot{\rho}^2(1 + \frac{\rho^2}{L^2}) + \frac{l_z^2}{2m\rho^2} , \quad V = \frac{1}{2}m\omega_0^2\rho^2(1 + \frac{1}{4}\frac{\rho^2}{L^2}) , \quad (3.9)$$

where $\omega_0^2 = g/L$. With two degrees of freedom and two constants of the motion (l_z and the energy), the system is integrable.

If the quartic terms are neglected in (3.9) by assuming $L \rightarrow \infty$, the (x, y) or (ρ, ϕ) motion is that of a 2D isotropic harmonic oscillator with frequency

ω_0 . In particular, an elliptical orbit, with semiaxes a and b , is fixed in space, and the motion is periodic. If the quartic terms are not negligible, the oscillator frequency shifts from ω_0 to a lower value ω , as in the plane pendulum case, where

$$\omega = \omega_0 \left(1 - \frac{a^2 + b^2}{16L^2} \right) , \quad (3.10)$$

and the orbit precesses in the prograde sense with frequency Ω , where

$$\Omega = \frac{3}{8} \frac{ab}{L^2} \omega_0 . \quad (3.11)$$

The motion is now in general quasiperiodic. The precession is one of the major sources of error to be eliminated in the design of Foucault pendula [33]. (The Foucault pendulum is discussed in Sec. 5.2.2.) Because the precession rate is $O(L^{-2})$, this precession is particularly relevant for recently developed short Foucault pendula, which are less than 1 m in length. Equations (3.10) and (3.11) have been derived by perturbation theory [34].

To derive (3.10) and (3.11) variationally [13], we first choose a trial trajectory; we take an ellipse which is closed in the coordinate system x', y' which rotates with respect to the fixed x, y system with the (unknown) precession frequency Ω . In the x', y' system we therefore have $x' = a \cos \omega t$ and $y' = b \sin \omega t$, where ω may differ from ω_0 . Because the x', y' and x, y systems differ by a rotation, we have $\rho^2 = x^2 + y^2 = x'^2 + y'^2$, and therefore our trial trajectory in the fixed coordinate system is

$$\rho^2 = a^2 \cos^2 \omega t + b^2 \sin^2 \omega t . \quad (3.12)$$

To apply the Maupertuis principles, we calculate the action W and mean energy $\bar{E}$ over one period $T = 2\pi/\omega$ of the rotating ellipse,

$$W = T \langle 2K \rangle, \quad \bar{E} = \langle K + V \rangle , \quad (3.13)$$

where $\langle \dots \rangle$ denotes a time average over one period, and K and V are given by (3.9). The averages of the terms in (3.9) are easily calculated using (3.12), which gives

$$\langle \rho^2 \rangle = \frac{1}{2} (a^2 + b^2) , \quad \langle \dot{\rho}^2 \rangle = \frac{\omega^2}{2} (a - b)^2 , \quad \left\langle \frac{1}{\rho^2} \right\rangle = \frac{1}{ab} ,$$

(3.14)

$$\langle \rho^2 \dot{\rho}^2 \rangle = \frac{\omega^2}{8}(a^2 - b^2)^2, \quad \langle \rho^4 \rangle = \frac{3}{8}(a^2 - b^2)^2 + a^2 b^2.$$

The precession frequency Ω can be related to the angular momentum using $\omega + \Omega \equiv \langle \dot{\phi} \rangle = \langle l_z / m \rho^2 \rangle$, and $\langle 1/\rho^2 \rangle = 1/ab$ from (3.14), giving

$$l_z = mab(\omega + \Omega) = mab\omega \left(1 + \frac{\Omega}{\omega}\right). \quad (3.15)$$




Figure 1: The precessing orbit. We choose positions 1 and 2 as the fixed end positions to be used in the action principle. (figure adapted from ref.[4])

As initial and final positions 1 and 2 (see Fig.1), we take two successive aphelia at $x' = a$, i.e. $(\rho_1, \phi_1) = (a, 0)$ and $(\rho_2, \phi_2) = (a, \phi_2)$, where $\phi_2 = (\omega + \Omega)T = 2\pi(1 + \Omega/\omega)$. The positions 1 and 2 and hence a and Ω/ω are thus fixed, which leaves b and ω available as variational parameters. Note that (3.15) then implies that l_z is varied with the trial trajectory (although it is a constant of the motion on the individual trial trajectories). We find it easiest to use the unconstrained version of the Maupertuis principles [i.e. the UMP (2.10)], $\delta \bar{E} = T^{-1} \delta W$, where $T = 2\pi/\omega$ is the time to reach position 2 from position 1. Thus we use

$$\frac{\partial}{\partial \omega} \bar{E} = \left(\frac{\omega}{2\pi}\right) \frac{\partial}{\partial \omega} W, \quad \frac{\partial}{\partial b} \bar{E} = \left(\frac{\omega}{2\pi}\right) \frac{\partial}{\partial b} W, \quad (3.16)$$

where, choosing units such that $m = 1$, $L = 1$, $\omega_0 = 1$, we have from (3.13) to (3.15)

$$\begin{aligned}\bar{E} = & \frac{\omega^2}{4}(a-b)^2 + \frac{\omega^2}{16}(a^2-b^2)^2 + \frac{\omega^2}{2}\left(1 + \frac{\Omega}{\omega}\right)^2 ab \\ & + \frac{1}{4}(a^2+b^2) + \frac{3}{64}(a^2-b^2)^2 + \frac{1}{8}a^2b^2, \end{aligned} \quad (3.17)$$

and

$$\frac{W}{4\pi} = \frac{\omega}{4}(a-b)^2 + \frac{\omega}{16}(a^2-b^2)^2 + \frac{\omega}{2}\left(1 + \frac{\Omega}{\omega}\right)^2 ab. \quad (3.18)$$

The first of the variational equations (3.16) is satisfied identically by (3.17) and (3.18), and gives no useful information. From the second, setting $\omega = \omega_0 + \Delta = 1 + \Delta$, where Δ is the frequency shift, and retaining terms only to second order in a and b , i.e., $O(L^{-2})$ in general units, we get the condition

$$\frac{a}{b}\Omega + \Delta = \frac{5}{16}a^2 - \frac{1}{16}b^2. \quad (3.19)$$

When $a = 0$, (3.19) gives $\Delta = -b^2/16$ as it should [recall the plane pendulum result [12, 13]]. By symmetry, Δ must reduce to $-a^2/16$ when $b = 0$. This requires that $\Omega = (3/8)ab$ in (3.19), and from this result we then get $\Delta = -(a^2 + b^2)/16$ from (3.19). When we restore the dimensional quantities ω_0 and L , these results agree with (3.10) and (3.11), so that with a harmonic oscillator trial trajectory, the variational method gives results correct to $O(L^{-2})$.

Hamilton's Principle can also be applied to this problem. Here, we need the Hamilton action $S = \int_0^T L dt = T\langle K - V \rangle$, where T (and hence ω) is now fixed, in addition to the positions 1 and 2 (and hence a and Ω/ω). This leaves b available as a variational parameter, and setting $\partial S/\partial b = 0$ leads to the same results as above.

4 More Complicated Examples. Research Problems

More complicated examples are easy to find, and are especially interesting in multidimensional systems which exhibit chaotic motion. Such systems are nonintegrable, and have fewer constants of the motion than the number

of degrees of freedom. We illustrate the use of classical and semiclassical variational principles together with the application of quantum variational principles to excited states.

4.1 The x^2y^2 oscillator

The 2D x^2y^2 oscillator is a simple example of this class, which has been studied in the literature for some time (see, e.g. [35, 36]). There is only one constant of the motion, the energy. Classically, most trajectories in this system are chaotic [37]. Fig.2 shows the contours of the x^2y^2 potential, and Fig.3 shows a typical (chaotic) classical trajectory.




Figure 2: Contours of the potential $V(x, y) = x^2y^2/2$. The darkened contour corresponds to $V(x, y) = 1$. (from ref.[35])

Refs.[12, 13] give classical and semiclassical discussions of the motion using the RMP. Here we discuss the quantum states using variational methods for the excited states, as done above for a simpler example. Comparison with the semiclassical results is given later. The Hamiltonian is

$$H = K_x + K_y + V(x, y) = \frac{p_x^2 + p_y^2}{2} + \frac{x^2y^2}{2}. \quad (4.1)$$




Figure 3: A typical trajectory of the Hamiltonian (4.1). The motion starts near the origin, then proceeds into the right-hand arm of the potential along the positive x -axis. The trajectory eventually reverses its x -direction and returns to the vicinity of the origin before entering another arm of the potential. (from ref.[35])

We choose units where $m = 1$ and $C = 1$, where $Cx^2y^2/2$ is the potential. Below we also choose $\hbar = 1$. We take as an approximation a separable trial wave-function $\psi_{n_x, n_y}(x, y)$

$$\psi_{n_x, n_y}(x, y) = \psi_{n_x}(\omega_x; x)\psi_{n_y}(\omega_y; y), \quad (4.2)$$

where the wave-function $\psi_{n_x}(\omega_x; x)$ is an eigenstate of a harmonic oscillator with frequency ω_x :

$$\frac{p_x^2 + \omega_x^2 x^2}{2} \psi_{n_x}(\omega_x; x) = (n_x + \frac{1}{2}) \omega_x \psi_{n_x}(\omega_x; x). \quad (4.3)$$

The frequencies ω_x and ω_y are variational parameters, which will differ from state to state, and as a result the set (4.2) is nonorthogonal. The computation of matrix elements and expectation values with the wave functions (4.3) is rather easy,

$$\langle n|K|n \rangle = \frac{1}{2}(n + \frac{1}{2})\omega, \quad \langle n|x^2|n \rangle = \frac{2n+1}{2\omega}, \quad (4.4)$$

and therefore the expectation value of the Hamiltonian (4.1) with the wavefunctions (4.2) is

$$\langle H \rangle = \langle n_x, n_y | H | n_x, n_y \rangle = \frac{1}{2}[(n_x + \frac{1}{2})\omega_x + (n_y + \frac{1}{2})\omega_y + \frac{(n_x + \frac{1}{2})(n_y + \frac{1}{2})}{\omega_x \omega_y}]. \quad (4.5)$$

We then find the best frequencies ω_x and ω_y by setting to zero the partial derivatives of $\langle H \rangle$ with respect to the two ω 's to obtain the frequency conditions

$$\omega_x^2 \omega_y = (n_y + \frac{1}{2}), \quad \omega_x \omega_y^2 = (n_x + \frac{1}{2}), \quad (4.6)$$

from which we obtain

$$(n_x + \frac{1}{2})\omega_x = (n_y + \frac{1}{2})\omega_y. \quad (4.7)$$

Eqs. (4.7) and (4.3) show that at the variational minimum the ratio of the two frequencies is rational and the mean energy of the x -oscillator is the same as the mean energy of the y -oscillator. It follows immediately that the mean kinetic energies of the two oscillators are also equal to each other at the variational minimum: $\langle K_x \rangle = \langle K_y \rangle$. From (4.6) we also find

$$\omega_x \omega_y = [(n_x + \frac{1}{2})(n_y + \frac{1}{2})]^{1/3}, \quad (4.8)$$

which gives the separate frequencies

$$\omega_x = (n_x + \frac{1}{2})^{-1/3}(n_y + \frac{1}{2})^{2/3}, \quad \omega_y = (n_x + \frac{1}{2})^{2/3}(n_y + \frac{1}{2})^{-1/3}, \quad (4.9)$$

and the variational estimate for the energy:

$$E_{n_x, n_y} = \langle n_x, n_y | H | n_x, n_y \rangle = \frac{3}{2}[(n_x + \frac{1}{2})(n_y + \frac{1}{2})]^{2/3}. \quad (4.10)$$

Therefore the variational states are labelled by the two integers n_x and n_y and the energy at high quantum numbers increases as the two thirds power of their product.

The estimate (4.10) can also be obtained semiclassically by using the RMP to find $\bar{E}(W_x, W_y)$, and then quantizing the actions W_i with the EBK relation (2.11); see ref.[13] for a detailed discussion. It is to be noted that this method yields the quite reasonable first approximation (4.10) for E_{n_x, n_y} despite the fact that strictly speaking the good actions W_x and W_y do not exist for the $x^2 y^2$ oscillator. In essence, one is approximating a chaotic trajectory by a quasiperiodic one, and the RMP picks out the best such trial trajectory [38].

The estimate (4.10) differs only in the leading coefficient, $3/2$ vs. 1.405 , from a semiclassical adiabatic formula (SCA) obtained by Martens et al [35]. The difference between the two estimates is about 6.5%. As noted by Martens et al [35] formula (4.10) gives rise to degeneracies. For example $(n_x, n_y) = (7,0), (0,7), (2,1), (1,2)$ give the same energies - a four-fold degeneracy. However, numerical estimates of the first 50 energy levels, obtained in [35], do not show such high degeneracies. Also, because the symmetry group of the Hamiltonian C_{4v} (a fourfold axis of rotation, and four reflection planes) has only one- and two-dimensional irreducible representations [39], we expect at most two-fold degeneracy. We shall see that the basis (4.2) permits an evaluation of the splitting of these spurious degeneracies. The pattern of degeneracies is easy to understand if the bracket is rewritten

$$[(2n_x + 1)(2n_y + 1)/4] = N/4,$$

where the number N (the “principal” quantum number), is an odd integer. The degeneracy pattern is determined by the decomposition of N into prime factors. For example $(n_x, n_y) = (2,1)$ corresponds to $N = 15$ which can be factored as $15 \cdot 1, 1 \cdot 15, 3 \cdot 5, 5 \cdot 3$ and these products correspond to $(7,0)$ etc. If N is a prime number there are only two states, corresponding to $N \cdot 1$ and $1 \cdot N$, while every nontrivial factorization of a nonprime N allows further states. The integer N belongs to one of two possible classes: $N = 4k + 1$ or $N = 4k - 1$. For the case $4k - 1$ (e.g. $N = 15$) one of n_x, n_y is even and the other is odd. This class of states EO and OE (of [35]) corresponds to the two dimensional irreducible representation of C_{4v} . For $N = 4k + 1$ (e.g. $N = 21$) the two quantum numbers (n_x, n_y) are both even or both odd (e.g. $(n_x, n_y) = (10,0)$ or $(0,10)$ or $(1,3)$ or $(3,1)$ for $N = 21$, which is therefore four-fold degenerate). This degeneracy is split into states of symmetry EEE, EEO, OOE and OOO, in the notation of Martens et al [35] for the one dimensional irreducible representations. In this notation the first two letters give respectively the parity (Even or Odd) of n_x and n_y (or equivalently the

behaviour of the wave-function under the reflections ($x \rightarrow -x$, $y \rightarrow -y$), while the third letter stands for the behavior under the interchange of x with y . We discuss later the splitting of degeneracies resulting from the expression (4.10). At this point we wish only to note that the degeneracy grows very slowly with N , like $\ln \ln N$, and even for N quite large where quasiclassical behaviour is expected (say $N \sim 105$) the typical number of degenerate states is quite modest, 4 - 6.

The main points of discussion arising from the result (4.10) are the splitting of the degenerate levels, perturbative corrections to the variational energies, and the density of states implied by this formula. For the first fifty low lying states one has the accurate numerical estimates of Martens et al [35] with which to compare.

4.1.1 Perturbative Shift of Variational Results

We can improve the variational result (4.10) if we rewrite the Hamiltonian H (4.1) in the form

$$H = H_0 + V = K + \frac{\omega_x^2 x^2 + \omega_y^2 y^2}{2} + \frac{x^2 y^2 - \omega_x^2 x^2 - \omega_y^2 y^2}{2}. \quad (4.11)$$

Here, the perturbation V is the last term above, and its effect is already included in first order, in the variational estimate (4.10). Therefore the first correction comes in second order in V . We must also remember that the frequencies (ω_x, ω_y) are appropriate to a fixed pair of quantum numbers (n_x, n_y) , and change if we go to a different pair. The simplest way to do perturbation theory in these circumstances is to estimate corrections to each state (n_x, n_y) by considering a set of orthogonal intermediate states (n'_x, n'_y) all with the same frequency (ω_x, ω_y) appropriate to the state (n_x, n_y) which we consider. Therefore we take the second order correction

$$\Delta E_{n_x, n_y}^{(2)} = - \sum \frac{|\langle n'_x, n'_y | V | n_x, n_y \rangle|^2}{[(n'_x - n_x)\omega_x + (n'_y - n_y)\omega_y]}. \quad (4.12)$$

Due to the form of V (quadratic in the coordinates x, y), we must have $n'_x = n_x$ or $n'_x = n_x + 2$ or $n'_x = n_x - 2$ for nonzero contributions and similarly for n'_y . It is an interesting property of the variational solution that only intermediate states with both n'_x different from n_x and n'_y different from n_y give nonvanishing contributions, because of the frequency conditions. This

can be seen from an evaluation of matrix elements with only one quantum number different, for example $n'_x \neq n_x$, but $n'_y = n_y$. Then one finds

$$\langle n'_x, n_y | V | n_x, n_y \rangle = \langle n'_x | \frac{x^2}{2} | n_x \rangle [\langle n_y | y^2 | n_y \rangle - \omega_x^2], \quad (4.12')$$

and the bracket on the right hand side vanishes because of (4.6). Only the perturbation $x^2 y^2 / 2$ contributes to nonvanishing matrix elements. There are only four intermediate states which contribute: $(+, +)$, $(+, -)$, $(-, +)$, $(-, -)$, where $(+, -) = (n_x + 2, n_y - 2)$, etc. Therefore (4.12) becomes after a little computation

$$\Delta E_{n_x, n_y}^{(2)} = -\frac{1}{24} E_{n_x, n_y}^{(0)}, \quad (4.13)$$

and

$$E_{n_x, n_y}^{(2)} = \frac{23}{16} [(n_x + 1/2)(n_y + 1/2)]^{2/3}, \quad (4.14)$$

where $\Delta E^{(2)}$ denotes a second-order shift and $E^{(2)}$ denotes the energy up to the second-order.

Therefore, in second-order perturbation theory, the coefficient $3/2$ is diminished by about 4% to 1.437, closer to the adiabatic result 1.405 of Martens et al [35]. The remaining difference is about 2.5%. The degeneracies of (4.14) are still the same as those of (4.10). If we apply perturbation theory to the ground state $(0, 0)$ we find

$$E_{00} = E_{00}^{(0)} + \Delta E_{00}^{(2)} + \Delta E_{00}^{(3)} + \Delta E_{00}^{(4)} = 0.5953 - 0.0248 + 0.0496 - 0.1465 + \dots \quad (4.15)$$

The sequence (4.15) indicates that the perturbation expansion does not converge, and it is a good idea to stop at second order. The accurate numerical result for E_{00} (from [35]) is 0.5541, which shows that stopping at second order gives an error of about 3% for the ground state.

Some examples of the estimates $E^{(0)}$, $E^{(2)}$, and E_{SCA} are given in Table 1, and compared with numerical estimates from [35], for a set of states which are also relevant to the next section.

Table 1

N	(n_x, n_y)	E_{exact}	$E_{var}^{(0)}$	$E^{(2)}$	E_{split}	E_{SCA}
1	(0,0)	0.5541	0.595	0.570	<i>na</i>	0.5577
5	(2,0)	1.7575	1.7406	1.6681	1.7471	1.6308
9	(4,0)	2.4920	2.5756	2.4683	2.5377	2.4132
13	(6,0)	3.1180	3.2911	3.1540	3.2171	3.0836
17	(8,0)	3.6909	3.9357	3.7717	3.8309	3.6874
21	(10,0)	4.2504	4.5310	4.3422	4.3973	4.2453
25	(12,0)	5.0490	5.0895	4.8775	4.9297	4.7685
29	(14,0)	5.3327	5.6189	5.3847	5.4347	5.2645

EEE states at low excitation. The exact values and E_{SCA} are taken from [35]; the variational estimates and second order results correspond to formulae (4.10) and (4.14), while the E_{split} are obtained following section 4.1.2

4.1.2 Splitting of Degenerate States

The basis (4.2) is also quite convenient to obtain the splitting of degenerate states. We shall deal with the simplest example with a degeneracy: the pair of states (2, 0) and (0, 2). Both these states have the same energy $(3/2)(5/4)^{2/3}$ in the variational approximation. The wave-functions of the two states are nonorthogonal, because the frequencies in the x and y oscillators differ. We define the overlap integral Δ

$$\Delta = \langle 20 | 02 \rangle = \left| \int dx \psi_2(\omega_2; x) \psi_0(\omega_0; x) \right|^2 = \frac{4}{27} \left(\frac{5}{4} \right)^{1/2}, \quad (4.16)$$

where the two wave functions have different frequencies ($\omega_2 = (1/10)^{1/3}$ and $\omega_0 = (25/2)^{1/3}$) as denoted. The splitting of two (or more) nonorthogonal degenerate states is well known; a simple example is the Heitler-London treatment of the hydrogen molecule. One has to take the sum or the difference of the two nonorthogonal states, e.g. $\Psi_+ = N_+[\psi_{02} + \psi_{20}]$, and similarly for Ψ_- . The eigenvalues have the form

$$E_+ = (H_{AA} + H_{AB}) / (1 + \Delta),$$

and

$$E_- = (H_{AA} - H_{AB}) / (1 - \Delta), \quad (4.17)$$

where here $|A\rangle \equiv |2, 0\rangle$ and $|B\rangle \equiv |0, 2\rangle$.

Thus in our case we need the off-diagonal matrix element of the Hamiltonian between the two states $(0, 2)$ and $(2, 0)$. This can be obtained from the overlap integral (4.16)

$$\langle 20|K_x + \frac{\omega_2^2 x^2}{2}|02\rangle = \frac{5}{2}\omega_2\Delta, \quad \langle 20|K_x + \frac{\omega_0^2 x^2}{2}|02\rangle = \frac{1}{2}\omega_0\Delta. \quad (4.18)$$

If we subtract the two equations (4.18) we obtain

$$\langle 20|(\omega_2^2 - \omega_0^2)\frac{x^2}{2}|02\rangle = (5\omega_2 - \omega_0)\Delta. \quad (4.19)$$

The right hand side of (4.19) vanishes because of the frequency condition (4.7) and therefore the matrix element of x^2 vanishes as well between the two variational states. The same is true for matrix elements of y^2 , and $x^2 y^2$. Therefore only the kinetic energy terms contribute to the off-diagonal matrix elements of the Hamiltonian. Using equations (4.18) we find

$$\langle 20|K_x|02\rangle = 12\frac{\omega_2^2\omega_0\Delta}{\omega_0^2 - \omega_2^2} = \frac{1}{2}\omega_0\Delta. \quad (4.20)$$

The same value is obtained for the matrix element of K_y , and therefore we find

$$\langle 20|H|02\rangle = \omega_0\Delta. \quad (4.21)$$

In our case $\omega_0 = (4/3)E_{02}$, and therefore we have for the two split eigenvalues

$$E_+ = E_{02}\frac{1 + (4/3)\Delta}{1 + \Delta} = 1.8230; \quad E_- = E_{02}\frac{1 - (4/3)\Delta}{1 - \Delta} = 1.6254. \quad (4.22)$$

Hence the two degenerate states at $E_{02} = 1.7406$ split into an EEE state at 1.823 and an EEO state at 1.625. We identify Ψ_+ as an EEE state because Ψ_+ has EEE symmetry, and similarly Ψ_- has EEO symmetry. When one compares with the numerical results of Martens et al [35], the estimates (4.22) are qualitatively correct; the error of E_+ is about +3.7% while the error of E_- is +6.4%. One can further apply the factor of 23/24, obtained in the previous section, to end with the estimates 1.7471 and 1.556. The result $E = 1.7471$ for the $(2,0)$ EEE state is given in Table I as E_{split} . Other low-lying EEE state energies are also given.

We end this brief excursion into the $x^2 y^2$ model with the conclusion that the quantum variational method for excited states [16] is particularly

well adapted for energy estimates in this system. We have used variation-perturbation theory here to demonstrate this method. The same results can also be obtained purely variationally, by using more elaborate trial states than (4.2); e.g., for an EEE state we would try (for n_x, n_y even)

$$\psi_{n_x, n_y}^{(+)}(x, y) = N_+ [\psi_{n_x}(\omega_x; x) \psi_{n_y}(\omega_y; y) + \psi_{n_y}(\omega_y; x) \psi_{n_x}(\omega_x; y)],$$

and similarly we would try $\psi_{n_x, n_y}^{(-)}(x, y)$ for the EEO states. We return to purely semiclassical variational methods in Sec. 4.3. For completeness, we note that classical variation-perturbation methods have also been devised [40].

4.2 The “Linear Baryon”

Another system very amenable to classical and quantum variational estimates is a system of three equal mass particles constrained to move along a straight line [41]. There are constant attractive forces between the particles, or a linear potential (as for 1D gravity), and the particles can move through each other. In suitable units the Hamiltonian can be expressed as

$$H = \frac{1}{2}[p_1^2 + p_2^2 + p_3^2] + \frac{1}{2}[|x_1 - x_2| + |x_2 - x_3| + |x_3 - x_1|], \quad (4.23)$$

where x_i are the positions and p_i the momenta. If we keep the centre of mass at the origin we have the constraints $x_1 + x_2 + x_3 = 0$ and $p_1 + p_2 + p_3 = 0$. We can then take internal Jacobi coordinates ρ, λ defined by $x_1 - x_2 = \rho\sqrt{2}$ and $x_1 + x_2 - 2x_3 = \lambda\sqrt{6}$, and the Hamiltonian becomes

$$H = \frac{1}{2}[p_\rho^2 + p_\lambda^2] + \frac{1}{2\sqrt{2}}[|\lambda\sqrt{3} - \rho| + |\lambda\sqrt{3} + \rho| + 2|\rho|]. \quad (4.24)$$

Taking now polar coordinates R, θ in the ρ, λ plane, where $\lambda = R \cos \theta$ and $\rho = R \sin \theta$, we obtain

$$H = \frac{1}{2}[p_R^2 + p_\theta^2/R^2] + \frac{1}{\sqrt{2}}R(|\sin \theta| + |\sin(\theta + 2\pi/3)| + |\sin(\theta - 2\pi/3)|). \quad (4.25)$$

The equipotentials of this Hamiltonian form regular hexagons. The six sides correspond to the six different permutations of the three particles in the

baryon; for example, one side corresponds to $x_1 < x_2 < x_3$, etc. The Hamiltonian (4.25) describes the motion of a particle inside this regular hexagon. This system has been studied in the literature [42] under the restriction to a single “wedge” (say $0 < \theta < \pi/3$) under the name of “wedge billiards in a gravitational field”. Recently also the relativistic generalization of the Hamiltonian (4.23) has been discussed [43].

The classical motion with the Hamiltonian (4.25) can be solved piecewise exactly. The trajectory consists of segments of parabolas in each wedge of the hexagon. One can visualize the system as a hexagonal “pit” or “funnel” in a gravitational field, with a point particle sliding inside the funnel. Since the system is not harmonic there are no normal modes, but an infinity of trajectories at each energy. Many of these can be simply described in terms of two prototypes having high or low “angular momentum” respectively. We use quotation marks since angular momentum is not conserved in a hexagonal potential. However the average angular momentum of specific orbits is still meaningful. The low “angular momentum” trajectory consists in a motion along a groove of the hexagonal funnel corresponding to $\theta = 0$, to the origin and continuing along $\theta = \pi$, and back. The high “angular momentum” trajectory is composed of segments of parabolas joined together to look like a circle. Each of these trajectories can sustain transverse oscillations to give rise to quasiperiodic motions. Examples are shown in references [41, 43]. Eventually for large transverse oscillations the two types of trajectories blur and chaotic motion ensues.

With only one constant of the motion (the energy) and two degrees of freedom, the system is nonintegrable, and possesses chaotic trajectories. These and the nonchaotic ones are piecewise parabolic, but it may be advantageous to represent them by simpler trial functions. For illustration we describe a low angular momentum trajectory $\theta = p_\theta = 0$ when (4.25) becomes

$$H = \frac{p_R^2}{2} + \frac{R\sqrt{3}}{2}. \quad (4.26)$$

Consider the trajectory with $E = 1$. The problem can be solved exactly to give a period (for the whole cycle) of $T_{exact} = 4.6188$, while a variational estimate with the RMP using a harmonic oscillator trial trajectory (with the same mean energy $\bar{E} = 1$ as the exact trajectory) gives $T_{variational} = 4.6526$, an overestimate of 0.73%.

Quantum variational methods are quite useful here too since we cannot solve the quantum problem exactly, either for the ground state or ex-




Figure 4: The quantum probability distributions for four excited states of the hexagon pit. The notation $n12m0$ means the state $n = 12, m = 0$. The color scale of relative probabilities is given at the bottom. The graphs have been rescaled to make the hexagon, the classical boundaries of the orbits for the given E , the same size.(from ref. [41])

cited states. To describe the symmetry species of the symmetry group C_{6v} of the quantum Hamiltonian we can use eigenfunctions of the 2D isotropic harmonic oscillator [44]. In polar coordinates these are products of an exponential $\exp(-\omega R^2/2)$ times a function of θ ($\sin(m\theta)$ or $\cos(m\theta)$) times $(\omega R^2)^{m/2}$ times a Laguerre polynomial in ωR^2 times a normalization factor. Although the azimuthal quantum number m is not conserved with the Hamiltonian (4.25) it is conserved modulo 6 due to the hexagonal symmetry of the problem. We can as a first approximation minimize the diagonal elements of the Hamiltonian (4.26) in this basis with respect to the frequency and obtain first order estimates for the energies of states which are labelled by a

principal quantum number n and m (modulo 6). For example the energy of the ground state $(n,m)=(0,0)$ is overestimated by about 0.39% in this basis while energy of the state $(n,m)=(15,0)$ is overestimated by about 3%. “Exact” energies can be obtained by diagonalizing the Hamiltonian in a large basis of harmonic oscillator states and minimizing eigenvalues with respect to ω . Another variational procedure uses products of Airy functions along the groove and perpendicular to it, and takes the scale factors of the two Airy functions as variational parameters. For the excited state $(15,0)$ the error is 0.014% (the “exact” energy is 16.34869 while the Airy approximation gives 16.3511). More examples can be found in Table 1 of reference [41]. The probability densities of a few states are shown in Fig. 4 where one can see one low angular momentum state $(12,0)$ and one high angular momentum state $(2,18)$ and two states which presumably correspond to chaotic trajectories.

4.3 He Atom Inside C_{70} Cage

Endohedral fullerene complexes such as He @ C_{70} have attracted much study in recent years, both experimental and theoretical (for reviews see [45, 46, 47, 48, 49]). Here X @ C_n denotes species X (atom, ion or molecule) encapsulated in a C_n cage [50]. Early theoretical work focussed on the prediction of infrared and Raman [53, 54] and neutron scattering [49] spectra due to the motion of X = He, Li^+ , Na^+ , K^+ and CO inside the near-spherical C_{60} cage. The predictions of [53] were subsequently confirmed experimentally [55, 56] for the infrared bands of $Li^+@C_{60}$. Confirmation of the predicted Raman bands [53] of $Li^+@C_{60}$ is tentative [56]. Later the theoretical studies were extended to study the dynamics and spectra of atoms, molecules and ions trapped in nonspherical cages, e.g. He @ C_{70} [57, 58, 59], Ne @ C_{70} [60], He @ C_{250} (nanotube) [61], Na^+ @ C_{120} (nanotube) [62], and even $(C_{60})_n$ @ C_∞ (nanotube), so-called nanopeapods [63]. Figs 5 and 6 show C_{60} , C_{70} and carbon nanotube cages.

The C_{60} cage, assumed to be rigid, presents a very nearly spherical potential [64] for an entrapped atom or ion X and the dynamics has been studied by various classical [66], semiclassical [48] and quantum [48] methods. The potential can be calculated by a variety of semiempirical and *ab initio* methods [47, 48, 66, 67]. The dynamics is usually quite regular [64] for X@ C_{60} . We discuss here instead the case He @ C_{70} , where the potential is highly nonspherical and gives rise to nonintegrable (chaotic) dynamics of the He atom




Figure 5: (a) The soccer ball-shaped molecule C_{60} (icosahedral symmetry group I_h); (b) The rugby ball-shaped molecule C_{70} (symmetry group D_{5d}). (from ref. [46])

[57, 58, 59]. We estimate the energy levels semiclassically [58, 59] using the UMP as described earlier, and compare with the “exact” results obtained by brute-force diagonalization of the Hamiltonian [57]. Classical simulations of the motion have also been carried out [57, 58, 59], and provide insight as to the degree of chaos in the motion of the He atom. The chaos is surprisingly small, given that the potential is strongly anharmonic. The small degree of chaos implies an approximate third constant of the motion [68], in addition to the energy and z -component of angular momentum (assuming the potential is axial).

The potential for He in a rigid C_{70} cage has been estimated semiempirically [57], and, to a good approximation, is axially symmetric around long z -axis in Fig 5b. With origin at the centre of the cage and x, y axes perpendicular to the symmetry z -axis, the potential has been fit to the form

$$V(x, y, z) = \frac{1}{2}k_1(x^2 + y^2) + \frac{1}{2}k_3z^2 + C(x^2 + y^2)z^2. \quad (4.27)$$

The first two terms describe a 3D anisotropic harmonic oscillator, with frequencies $\omega_x^{(0)} = \omega_y^{(0)} = (k_1/m)^{1/2} = 84.166 \text{ cm}^{-1}$ and $\omega_z^{(0)} = (k_3/m)^{1/2} = 21.934 \text{ cm}^{-1}$, where m is the He atom mass. We use the standard spectroscopic unit (cm^{-1}) for frequency and energy (see [58]). The last term $C(x^2 + y^2)z^2$ in (4.27) is a quartic perturbation, with $C = 1.738 \text{ cm}^{-1}$ when x, y, z are in units of $(\hbar/m\omega_x^{(0)})^{1/2}$. There are also smaller terms in (4.27) varying as $(x^2 + y^2)^2$, z^4 , etc, which we shall ignore.

Because of the axial symmetry we change to cylindrical polar coordinates




Figure 6: Carbon nanotube of diameter 0.7 nm. This is a quite narrow nanotube with the hemispherical endcaps derived from C_{60} . The molecule depicted is C_{250} ; actual nanotubes are much longer [65], up to $C_{1,000,000}$ and beyond. (from ref [61])

(ρ, ϕ, z) and write the Hamiltonian, with potential (4.27), as

$$H = \frac{1}{2}m(\dot{\rho}^2 + \dot{z}^2) + \frac{l_z^2}{2m\rho^2} + \frac{1}{2}m\omega_x^{(0)2}\rho^2 + \frac{1}{2}m\omega_z^{(0)2}z^2 + C\rho^2z^2, \quad (4.28)$$

where $\rho = (x^2 + y^2)^{1/2}$, and l_z is the z -component of the angular momentum, which is a constant of the motion. The cage has been assumed to be rigid and fixed in space. To apply the UMP (2.10), we proceed exactly as for the simpler spherical pendulum example of sec.3.2. When viewed down the z axis, the trajectory is roughly a precessing ellipse, and the z -motion is oscillatory. We therefore take as a trial trajectory

$$x' = A_x \cos \omega_x t, \quad y' = A_y \sin \omega_x t, \quad z' = z = A_z \cos \omega_z t, \quad (4.29)$$

where the x', y' axes precess with some (unknown) frequency Ω with respect to x, y , and where ω_x may differ from $\omega_x^{(0)}$, etc. Calculating $\bar{E}$ and W with the trial trajectory and applying the UMP gives to first order in C (see [58] for the details) $\Omega = 0$, the shifted frequencies

$$\omega_x = \omega_x^{(0)} + \frac{CW_z}{2\pi m^2 \omega_x^{(0)} \omega_z^{(0)}}, \quad \omega_z = \omega_z^{(0)} + \frac{C(W_x + W_y)}{2\pi m^2 \omega_x^{(0)} \omega_z^{(0)}}, \quad (4.30)$$

and also, to first-order in C , $\bar{E}$ as a function of the partial actions $I_i = W_i/2\pi$

$$\bar{E}(I_x, I_y, I_z) = (I_x + I_y)\omega_x^{(0)} + I_z\omega_z^{(0)} + \frac{C(I_x + I_y)I_z}{m^2\omega_x^{(0)}\omega_z^{(0)}}. \quad (4.31)$$

The terms of second-order in C in (4.30) and (4.31) are also derived in [58], and it is also found that $\Omega = 0$ to second order. The simulations show that Ω is small but nonzero in general, and to obtain $\Omega \neq 0$ from the UMP requires an improved trial trajectory over (4.29), e.g. one involving higher harmonics of ω_x and ω_y [59]. We shall ignore this small effect here.

To estimate the energy levels semiclassically from (4.31) is very simple: according to the EBK quantization rule (2.11), we replace I_x by $(n_x + \frac{1}{2})\hbar$, etc. This gives

$$E_{n_x, n_y, n_z} = (n_x + n_y + 1)\hbar\omega_x^{(0)} + (n_z + \frac{1}{2})\hbar\omega_z^{(0)} + \frac{C(n_x + n_y + 1)(n_z + \frac{1}{2})\hbar^2}{m^2\omega_x^{(0)}\omega_z^{(0)}} + O(C^2) \quad (4.32)$$

where the $O(C^2)$ and $O(C^3)$ terms are written down in ref. [58]. We compare the semiclassical energy levels (4.32) (including the $O(C^2)$ and $O(C^3)$ terms) with the fully quantum results; the lowest 613 levels, up to the energy of 1000 cm^{-1} are obtained by diagonalizing the Hamiltonian in a harmonic oscillator basis in [58]. The harmonic oscillator chosen has a potential given by the harmonic part of (4.28). The semiclassical and fully quantum results agree to better than 1%. We illustrate the good agreement in Fig 7, which shows the levels up to 500 cm^{-1} , organized according to m , the quantum number corresponding to l_z .

For an isotropic 2D harmonic oscillator, the energy depends only on $N_{xy} = n_x + n_y$. For $N_{xy} = 2$, for example, we can have $(n_x, n_y) = (2, 0), (1, 1), (0, 2)$, which are degenerate. The m -values allowed for $N_{xy} = 2$ are $m = -2, 0, +2$. For general N_{xy} , we have $|m| = N_{xy}, N_{xy} - 2, N_{xy} - 4, \dots, 1$ or 0 . For our oscillator (4.28) the different $|m|$ states should in reality have different energies; this is found in the fully quantum calculation, but not in the semiclassical one where the energy depends only on $N_{xy} = n_x + n_y$. Table 2 illustrates more clearly the slight $|m|$ splittings found quantum mechanically, but not semiclassically.

To obtain the $|m|$ splittings semiclassically requires an improved trial trajectory over (4.29), one that gives a nonvanishing precession frequency Ω . This is discussed in ref [59]. The simple improvement in the trial trajectory




Figure 7: Comparison of energy levels calculated via quantum mechanical methods (black) and a semiclassical variational technique (grey).(from ref.[58])

mentioned earlier (including some higher harmonics) does indeed give a non-vanishing Ω and hence $|m|$ splitting. However the trial trajectories employed thus far are unable to give accurate splittings, and hence require further improvement.

Table 2

n_z	$N_{xy} = n_x + n_y$	m	$E_{quant}(cm^{-1})$	$E_{semiclass}(cm^{-1})$
0	2	2	262.818	263.466
0	2	0	262.821	263.466
1	2	2	284.696	285.403
1	2	0	284.699	285.403
2	2	2	306.574	307.340
2	2	0	306.577	307.340

Comparison of selected energy levels calculated via quantum and semiclassical methods demonstrating the failure of the simple variational calcula-

tion to produce level splitting observed in the quantum results.(from ref.[58])

5 Nonconservative Systems

In this section we consider nonconservative mechanical systems. For example consider a system of particles moving in the time-dependent external field, or a system whose parameters depend explicitly on time. Other examples include time-dependent constraints, e.g. a bead on a rotating hoop whose angular velocity is a prescribed function of time, and variable mass systems (for which $L \neq K - V$ in general [69]). Such systems are not closed and the energy is not conserved. The evolution of a nonconservative system is completely determined by a time-dependent Lagrangian $L(q, \dot{q}, t)$ or by a time-dependent Hamiltonian $H(q, p, t)$. (We exclude dissipative systems from our discussion. We do note that Lagrangians and Hamiltonians have been found for such systems in some cases [70], so that the action principles discussed here would apply in these cases. Alternatively [5], dissipative terms can be added to the Lagrange equations of motion or to the Hamilton principle.) H and L are related to each other by the Legendre transformation (see eq. (2.5))

$$H(q, p, t) = \sum_n p_n \dot{q}_n - L(q, \dot{q}, t). \quad (5.1)$$

Locally these lead to the usual Euler-Lagrange or Hamilton equations of motion that govern the dynamics of the system.

5.1 Action Principles

We consider the global description of nonconservative systems. It is well known that the Hamilton Principle

$$(\delta S)_T = 0 \quad (5.2)$$

is applicable for nonconservative systems as well as for conservative ones. Here subscript T reminds us that variations near the actual path are constrained by fixing the travel time T from the initial point $A \equiv (q_A, 0)$ to the final point $B \equiv (q_B, T)$. Again we have chosen $t_A = 0$ and $t_B = T$, and, as before, it is understood here and below that the end-positions q_A and q_B are also fixed.

As before, we can relax the constraint of fixed T and rewrite (5.2)[12, 71] as an unconstrained Hamilton Principle (UHP)

$$\delta S + E(T)\delta T = 0, \quad (5.3)$$

where the Lagrange multiplier $E(T)$ is the true value of the energy at the time of arrival at the final point B . (Elsewhere in this paper $E(T)$ is denoted variously as $E(t_B)$, E_B or H_B ; see notes [25] and [73]). We assume that all trial trajectories start at the same time $t = 0$ and also have the same end positions q_A and q_B .

Using eq.(2.7), which is derived from (5.1) as before, we transform the UHP (5.3) to the form

$$T\delta\bar{E} - \delta W - [E(T) - \bar{E}(T)]\delta T = 0. \quad (5.4)$$

This is the unconstrained Maupertuis Principle (UMP) for nonconservative systems, which seems not to have been given in the literature explicitly (it is implicit in an Appendix in [12] - see also [73]). It relates three variations - the variation of mean energy $\bar{E}$, the variation of the action W and the variation of the travel time T . The Lagrange multipliers are the true travel time T and the difference of energy and mean energy of the true trajectory at the final point, $[E(T) - \bar{E}(T)]$. The latter can be rewritten as

$$E(T) - \bar{E}(T) = \frac{1}{T} \int_0^T dt [t \frac{\partial H}{\partial t}] = \langle t \frac{\partial H}{\partial t} \rangle, \quad (5.5)$$

where the bar and brackets $\langle \dots \rangle$ both denote a time average over the interval $(0, T)$. The generalized UMP (5.4) can also be derived by traditional variational procedures ([12], eq.(A.6)).

One can derive constrained principles from (5.4) as before, e.g. $(\delta\bar{E})_{W,T} = 0$, but these would be complicated to use in practice because of the severe constraints on the trial trajectories. The version $(\delta W)_{\bar{E},T} = 0$ is closest to the original Maupertuis Principle. The constrained principle

$$\left(\delta [W + T \langle t \frac{\partial H}{\partial t} \rangle] \right)_{E(0), E(T)} = 0$$

is mentioned by Whittaker [75], and is easily derived from (5.4) and (5.5). But since (5.4) is also valid without fixing $E(0)$ and $E(T)$, (5.4) is more convenient.

Formally the unconstrained Maupertuis principle (5.4) is equivalent to the Hamilton principle, but it also appears too complicated to be useful in practice. This is not quite true and we demonstrate how the UMP can be used for adiabatic perturbations.

5.2 Adiabatic Invariants and the Hannay Angle

For simplicity we start with a one-dimensional periodic system that depends on a parameter λ . Let λ vary with time adiabatically, i.e. the change $\Delta\lambda \ll \lambda$ during a period of oscillation T . It is well known that in this case the energy E as well as the period of oscillation T depend slightly on time, but the action I over a cycle, where

$$I = \frac{1}{2\pi} W(\text{cycle}) = \frac{1}{2\pi} \oint p dq, \quad (5.6)$$

remains constant to an excellent approximation (adiabatic invariant) when the parameter λ varies.

Consider what kind of information can be extracted from the UMP (5.4). Both the energy E and the period T of an actual motion are functions of the action I (recall the action-angle variables): $E = E(I)$, $T = T(I)$. Consider two actual trajectories with actions I and $I + dI$ as two trial trajectories for the UMP (5.4). Then the variational equation can be rewritten as a differential equation

$$\frac{d\bar{E}}{dI} - \frac{2\pi}{T} = \langle t \frac{\partial E}{\partial t} \rangle > \frac{1}{T} \frac{dT}{dI}. \quad (5.7)$$

It is more convenient to rewrite this in terms of the frequency $\omega(I) = 2\pi/T(I)$,

$$\omega(I) = \frac{d\bar{E}}{dI} + \langle t \frac{\partial E}{\partial t} \rangle > \frac{1}{\omega(I)} \frac{d\omega(I)}{dI}. \quad (5.8)$$

This equation generalizes the standard relation [5] for integrable conservative systems

$$\omega(I) = \frac{dE}{dI}$$

to integrable nonconservative systems.

Up to this point (5.7) and (5.8) are general and valid for any nonconservative system for which a good action can be defined. Consider now the

adiabatic regime where the variation of the parameter $\lambda(t)$ over the short period T , $\Delta\lambda = \lambda(T) - \lambda(0)$, is much less than the parameter itself: $\Delta\lambda \ll \lambda(0)$. In this case

$$\langle t \frac{\partial E(I, \lambda(t))}{\partial t} \rangle = \langle t \frac{\partial E(I, \lambda)}{\partial \lambda} \dot{\lambda} \rangle = \dot{\lambda} \langle t \frac{\partial E(I, \lambda)}{\partial \lambda} \rangle, \quad (5.9)$$

where the averaging $\langle t \partial E(I, \lambda) / \partial \lambda \rangle$ is to be performed at a fixed value of $\lambda \simeq \lambda(0)$ since λ varies little over a period T . The action I is conserved for any given parameter λ , and hence the derivative $\partial E(I, \lambda) / \partial \lambda$ is also a constant of the motion. As a result in this approximation we get

$$\langle t \frac{\partial E(I, \lambda)}{\partial \lambda} \rangle = \frac{T}{2} \frac{\partial E(I, \lambda)}{\partial \lambda} = \frac{\pi}{\omega} \frac{\partial E(I, \lambda)}{\partial \lambda}.$$

In the same approximation we have for $\bar{E} \equiv \langle E \rangle$ in (5.8)

$$\langle E(I, \lambda(t)) \rangle = E(I, \lambda(0)) + \frac{\pi}{\omega} \dot{\lambda} \frac{\partial E}{\partial \lambda}.$$

Thus to the first order approximation in $\dot{\lambda}$ we can rewrite (5.8) as

$$\omega(I) = \frac{dE(I, \lambda(0))}{dI} + [\pi \dot{\lambda}] \left[\frac{d}{dI} \left(\frac{1}{\omega^{(0)}} \frac{\partial E}{\partial \lambda} \right) - \frac{\partial E}{\partial \lambda} \frac{d}{dI} \left(\frac{1}{\omega^{(0)}} \right) \right],$$

or finally as

$$\omega(I) = \omega^{(0)} + \frac{\Delta\lambda}{2} \frac{\partial \omega^{(0)}}{\partial \lambda}. \quad (5.10)$$

Here $\omega^{(0)}(I, \lambda) = dE(I, \lambda(0)) / dI$ is the zero-order term, and $\Delta\lambda = \lambda(T) - \lambda(0) \simeq \dot{\lambda}(2\pi / \omega^{(0)})$ is the variation of the parameter λ during the given period. We get corrections (of the first-order in the nonadiabaticity parameter $\dot{\lambda}$) to the frequency of oscillation, for adiabatic systems. The result seems very simple. It appears that $\omega(I)$ is a continuous function of time t and we calculate some average frequency over the period. But actually $\omega(I)$ is not always a continuous function of time; it is a global parameter of a (slightly) aperiodic system. Thus (5.10) is nontrivial in general.

To illustrate these results it is instructive to consider a simple exactly solvable example. We consider the 1D motion of a perfectly elastic ball of mass m bouncing between two planes. We suppose that one plane is fixed and the other is slowly moving with constant velocity u . We suppose also

that the planes are heavy enough so that they are unaffected by collisions with the ball. The distance between the two planes $L(t) = L_0 + ut$ is the adiabatic parameter.

Let us define a cycle as the motion of the ball from the fixed plane to the moving plane, collision, and the motion of the ball in the backward direction up to the collision with the fixed plane. The n -th cycle can be described in the following way. At time $t = 0$ (beginning of the cycle) the ball has velocity v_n , and the distance between planes is L_n . At time $T_n(\rightarrow) = L_n/(v_n - u)$ the ball collides with the moving plane and after the collision it moves to the fixed plane with velocity $(v_n - 2u) = v_{n+1}$ (the collision with the fixed plane is elastic and the backward velocity is equal to the forward velocity after collision, i.e. v_{n+1}). The backward motion takes time $T_n(\leftarrow) = (L_n v_n)/(v_n - u)v_{n+1}$. The period of the whole cycle is $T_n = T_n(\rightarrow) + T_n(\leftarrow) = (2L_n)/v_{n+1}$. At time $t = T_n$ (the initial time for $(n+1)$ -th cycle) the distance between the two planes is equal to $L_{n+1} = L_n v_n/v_{n+1}$. This means that $L_n v_n = L_{n+1} v_{n+1}$ is an exact (adiabatic) invariant.

It easy to see that the action I is proportional to this invariant,

$$I_n = \frac{1}{2\pi} \oint p dq = \frac{m}{\pi} L_n v_n,$$

i.e. the action I is an exact (adiabatic) invariant for our special initial condition [74].

In terms of the action I and the adiabatic parameter L the complete set of relevant quantities is as follows:

$$\begin{aligned} \bar{E}_n &= \frac{1}{2m} \left(\frac{\pi I}{L_n} \right)^2 - u \left(\frac{\pi I}{L_n} \right), \\ E(T_n) &= \frac{1}{2m} \left[\left(\frac{\pi I}{L_n} \right) - 2mu \right]^2, \\ \omega_n(I) &= \frac{2\bar{E}_n}{I} = \frac{1}{2m} \left(\frac{\pi^2 I}{L_n^2} \right) - u \left(\frac{\pi}{L_n} \right), \end{aligned}$$

i.e. $\omega_n(I)$ is linear function of the nonadiabaticity parameter $\dot{L} = u$ and the approximate relation (5.10) must be exact here. Indeed, we have $\omega^{(0)} = \pi^2 I / 2m L_n^2$, $\Delta L = \pi u / \omega^{(0)}$ and

$$\omega^{(0)} + \frac{\Delta L}{2} \frac{\partial \omega^{(0)}}{\partial L} = \frac{1}{2m} \left(\frac{\pi^2 I}{L_n^2} \right) - u \left(\frac{\pi}{L_n} \right) = \omega(I).$$

Thus the approximate relation (5.10) is valid here. It is also not difficult to check after a little algebra that in the exact relation (5.8) all higher powers of u cancel each other and the linear approximation is exact.

If $u \equiv \dot{L}$ is not extremely small, I ceases to be an adiabatic invariant and the system becomes nonintegrable. If $L(t)$ is varied nonadiabatically and periodically around L_0 , the system (termed the Fermi model [76]) can show steady energy growth and chaotic motion.

The generalization of (5.10) to multidimensional integrable systems with actions $I_1, \dots, I_f$ is evident, as is the generalization to multiparameter systems, with parameters $\lambda_1, \lambda_2, \dots$.

The $O(\dot{\lambda})$ corrections in $\omega(I)$ have been derived by other methods [77], but in different forms from (5.10). If we assume $\lambda(t)$ is periodic, with a long period T_λ , the $O(\dot{\lambda})$ terms in $\omega(I)$, when integrated over $(0, T_\lambda)$, define the Hannay angle $\Delta\theta_H$, the classical analogue of the quantum Berry phase [77]. The Hannay angle is the extra (geometric) phase undergone by the system in one cycle of $\lambda(t)$, in addition to the more obvious dynamical phase θ_D obtained by integrating the instantaneous frequency $\omega(\lambda(t)) \equiv dE(I, \lambda(t))/dI$. In general one requires more than one λ -type parameter to obtain a nonvanishing result for $\Delta\theta_H$, but Hannay and Berry and coworkers give a few examples [77, 78, 79] where one parameter suffices. We discuss two examples with nonvanishing $\Delta\theta_H$.

5.2.1 Bead Sliding on Slowly Rotating Horizontal Ring

Consider a bead of mass m sliding without friction on a closed planar wire loop which is slowly rotating with angular velocity Ω in the horizontal plane. For a loop of arbitrary shape, Hannay and Berry [77] show that the Hannay angle $\Delta\theta_H$ defined above is given by $\Delta\theta_H = -8\pi^2 A/L^2$, where A is the area and L the perimeter of the loop. For a circular loop rotating about an axis perpendicular to the plane of the loop and through the center, $\Delta\theta_H$ is simply -2π , as is clear intuitively: the “starting line” for the particle moves ahead by 2π while the particle rapidly slides around the loop.

We consider a circular loop of radius a (Fig.8), but with the vertical axis of rotation through a point O on the wire instead of the center C of the loop, to make the answer less obvious. The adiabatic parameter here is the prescribed orientation of the loop $\lambda(t)$, where $\dot{\lambda}(t) \equiv \Omega(t)$ is small. We do not at this stage assume $\Omega(t)$ is constant.

Using the horizontal x, y lab-frame (Fig.8), we can easily write the La-




Figure 8: Circular wire loop rotating slowly in the horizontal x, y plane with angular velocity $\Omega = \dot{\lambda}$ about axis through O perpendicular to the x, y plane. At $t = 0$, the particle starts with $\theta = 0$ and the rotating x' axis (a diameter) coincides with the fixed axis x .

grangian, which is just the lab kinetic energy of the particle. Using θ in Fig.8 as the generalized coordinate we find

$$L = \frac{1}{2}ma^2 \left[\Omega^2 + (\Omega + \dot{\theta})^2 + 2\Omega(\Omega + \dot{\theta}) \cos \theta \right] . \quad (5.11)$$

From (5.11) we obtain the equation of motion

$$\ddot{\theta} + \dot{\Omega}(1 + \cos \theta) + \Omega^2 \sin \theta = 0 . \quad (5.12)$$

If the loop rotates uniformly ($\dot{\Omega} = 0$), (5.12) is just the pendulum equation, as one expects intuitively.

Because we are using a rotating coordinate system (θ is measured from x') L is not equal to $K - V \equiv K = \frac{1}{2}ma^2\dot{\theta}^2$ in this system, so that the Hamiltonian H is not equal to the energy ($= \frac{1}{2}ma^2\dot{\theta}^2$) in this system either. To find H , we first calculate the canonical momentum p_θ , where

$$p_\theta = \frac{\partial L}{\partial \dot{\theta}} = ma^2(\Omega + \dot{\theta} + \Omega \cos \theta) , \quad (5.13)$$

and then $H = p_\theta \dot{\theta} - L$ is

$$H = \frac{1}{2}ma^2\dot{\theta}^2 - ma^2\Omega^2(1 + \cos \theta) . \quad (5.14)$$

In terms of the canonical variables θ , p_θ the Hamiltonian H is

$$H = \frac{p_\theta^2}{2ma^2} - p_\theta\Omega(1 + \cos\theta) + O(\Omega^2) \quad , \quad (5.15)$$

where we drop $O(\Omega^2)$ terms since we assume $\Omega \ll \dot{\theta}$.

We cannot apply (5.10) as it stands, since $H = E$ was used in the derivation. The generalization to include cases where $H \neq E$ is

$$\omega(I, \lambda) = \frac{d\bar{H}}{dI} - (\pi\dot{\lambda}\frac{\partial\bar{H}}{\partial\lambda})\frac{d}{dI}(\frac{1}{\omega^{(0)}(I)}), \quad (5.16)$$

where $\bar{H}$ is the average of H over the fast motion (here θ). Alternatively, we can use (5.10), with now $\omega^{(0)} = dH(I, \lambda(0))/dI$. From (5.15) we have

$$\bar{H} \doteq \frac{p_\theta^2}{2ma^2} - p_\theta\Omega, \quad (5.17)$$

since θ varies rapidly and p_θ is nearly constant.

We now assume Ω is constant. For $\Omega = 0$, $\bar{H}$ is a function of the good action $I = p_\theta$ only (no θ dependence), and when $\Omega \neq 0$ we see p_θ remains a good action. Note that $\bar{H}$ is independent of angle λ , so that the second term in (5.16) will not contribute to $\omega(p_\theta) = d\bar{H}/dp_\theta$, which is

$$\omega(p_\theta) = \frac{p_\theta}{ma^2} - \Omega \quad . \quad (5.18)$$

The total phase undergone after a long return time $T_\Omega = 2\pi/\Omega$ is thus

$$\theta = (\frac{p_\theta}{ma^2})T_\Omega - 2\pi \quad . \quad (5.19)$$

The first term is the dynamical phase, the second is the Hannay (geometric) phase.

5.2.2 The Foucault Pendulum

For a Foucault pendulum the Hannay angle is the angle by which the plane of the pendulum vibration shifts in one day, i.e. $\Delta\phi_H = -2\pi \cos\theta$, where θ is the co-latitude of the point on earth where the pendulum is located. This is a clockwise rotation in the northern hemisphere. This result for $\Delta\phi_H$ has been derived by Berry [79] and others [80] by various arguments. There are also

some pre-Hannay purely geometric arguments for the Foucault pendulum precession [81]. Here we show the result follows simply from (5.16).

We consider a pendulum of length L and bob mass m oscillating at a point on the earth's surface with co-latitude θ . We use axes x, y, z attached to the earth, with x, y in the horizontal plane and z vertical. When at rest in the x, y, z system, the pendulum hangs along the z -axis, and when in motion the bob oscillates very nearly in the x, y plane. Neglecting the z motion and assuming $\Omega \equiv \Omega_E \cos \theta$ is small, where Ω_E is the earth's angular velocity, we easily find [82] the approximate horizontal equations of motion

$$\ddot{x} + \omega_0^2 x - 2\Omega \dot{y} = 0, \quad \ddot{y} + \omega_0^2 y + 2\Omega \dot{x} = 0, \quad (5.20)$$

where $\omega_0 = (g/L)^{1/2}$. We have kept the $O(\Omega)$ Coriolis terms, but dropped the $O(\Omega^2)$ centrifugal terms, since Ω is small compared to the pendulum frequency ω_0 . Here Ω is the local vertical component of the earth's angular velocity. We are interested in particular in the orbit precession rate predicted by (5.20).

It is easily checked that (5.20) follow from the Lagrangian

$$L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) - \frac{1}{2}m\omega_0^2(x^2 + y^2) + m\Omega(xy - yx), \quad (5.21)$$

and of course (5.21) can be derived directly [83].

Changing to polar coordinates ρ, ϕ via $x = \rho \cos \phi, y = \rho \sin \phi$ changes L to the form

$$L = \frac{1}{2}m\dot{\rho}^2 + \frac{1}{2}m\rho^2\dot{\phi}^2 - \frac{1}{2}m\omega_0^2\rho^2 + m\Omega\rho^2\dot{\phi}. \quad (5.22)$$

Two constants of the motion are obvious from (5.22): since L does not depend on ϕ , p_ϕ is a constant, where

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = m\rho^2\dot{\phi} + m\rho^2\Omega; \quad (5.23)$$

and because L does not depend on t explicitly, the Hamiltonian $H = p_\rho\dot{\rho} + p_\phi\dot{\phi} - L$ is a constant, where $p_\rho = \partial L/\partial \dot{\rho}$ and

$$H = \frac{1}{2}m\dot{\rho}^2 + \frac{1}{2}m\rho^2\dot{\phi}^2 + \frac{1}{2}m\omega_0^2\rho^2. \quad (5.24)$$

Two constants of the motion with two degrees of freedom imply that the system is integrable and that two good actions exist.

To express H in terms of the canonical variables $\rho, \phi, p_\rho, p_\phi$ we use (5.23) and

$$p_\rho = \frac{\partial L}{\partial \dot{\rho}} = m\dot{\rho} \quad (5.25)$$

to get

$$H = \frac{p_\rho^2}{2m} + \frac{p_\phi^2}{2m\rho^2} + \frac{1}{2}m\omega_0^2\rho^2 - \Omega p_\phi, \quad (5.26)$$

where again we drop an $O(\Omega^2)$ term. We wish to use (5.16) (not (5.10) here as $H \neq K + V$) to find the frequencies $\omega_\rho \equiv \omega(I_\rho, \lambda)$ and $\omega_\phi = \omega(I_\phi, \lambda)$, where λ is the orientational angle of the earth multiplied by $\cos\theta$. Note that for $\Omega = \dot{\lambda} = 0$, (5.26) is simply the Hamiltonian for an isotropic 2D harmonic oscillator. Calculation of the actions for the 2D oscillator, I_ρ and $I_\phi \equiv p_\phi$, is a standard exercise [84], and the Hamiltonian $H(\Omega = 0)$ in terms of these actions is

$$H(\Omega = 0) = 2I_\rho\omega_0 + |I_\phi|\omega_0. \quad (5.27)$$

The term $\Omega p_\phi \equiv \Omega I_\phi$ in (5.26) will not disturb the good actions, so that (5.26) becomes

$$H = 2I_\rho\omega_0 + |I_\phi|\omega_0 - I_\phi\Omega. \quad (5.28)$$

Application of (5.16) with $I = I_\rho$ and $I = I_\phi$ then gives

$$\omega_\rho = 2\omega_0, \quad \omega_\phi = \pm\omega_0 - \Omega, \quad (5.29)$$

where a positive (+) frequency in ω_ϕ corresponds to a counterclockwise rotation. Note that the Foucault precession frequency $\omega_F \equiv -\Omega$ is true for orbits of any shape, e.g. near linear (the usual case), elliptical, or circular. The factor of 2 in ω_ρ in (5.29) is clear intuitively: for $\Omega = 0$, in one circuit of an elliptical orbit, ρ goes through two cycles while ϕ goes through one. Computing the phase of the pendulum $\phi(T_E)$ after one complete rotation of the earth (period T_E) then gives

$$\phi(T_E) = \pm\omega_0 T_E - \Omega T_E = \pm\omega_0 T_E - 2\pi \cos\theta. \quad (5.30)$$

The term $\pm\omega_0 T_E$ is the dynamical phase ϕ_D and $-2\pi \cos\theta$ is the geometric Hannay angle $\Delta\phi_H$.

The examples in this section and the previous one led to a nonvanishing Hannay angle $\Delta\theta_H$. It is easy to show that the earlier example (particle in

a 1D box with a slowly moving wall) gives a vanishing result. The same is true for a particle in a slowly rotating 1D box (fixed walls). Golin [85] has proved that, for 1D systems with one adiabatic parameter λ , one always has $\Delta\theta_H = 0$ unless λ undergoes a rotational excursion through 2π . However the proof does not cover our rotating 1D box example (which has a hard wall potential), since Golin [85] assumes a smooth potential. The case of a slowly rotating 2D box of elliptical shape has been worked out by Koiller [80], and gives a nonvanishing $\Delta\theta_H$ depending in a complicated way on the area of the box.

5.2.3 Nonintegrable Systems

Nonintegrable systems have fewer than f independent constants of the motion or good actions, where f is the number of degrees of freedom. For such systems, at least some of the possible motions are chaotic, and for these at least some of the good action variables do not exist. Hence we have not been able to proceed rigorously beyond the general relation (5.4) for chaotic motions. (Despite the nonexistence of good actions I for nonintegrable systems, formal use of them and their adiabatic invariance have been used successfully in semiclassical quantization schemes; see sections 2.3, 3.1, 4.1, 4.3 and ref. [13] for a method based on the Maupertuis Principles for conservative systems, and see refs. [91, 92] for a method based on adiabatic switching.)

As stated, we do not have rigorous arguments to draw consequences from the UMP (5.4) for nonintegrable systems, but some heuristic arguments can be given. For example, by choosing $H(\lambda(t))$ to vary slowly and linearly with t , e.g. $H(\lambda(t)) = \lambda(t)H_0(q, p) = (1 + \dot{\lambda}t)H_0(q, p)$, where $\dot{\lambda}$ is small, we can derive the adiabatic invariance of the action I for a 1D periodic system (earlier we assumed that I is an adiabatic invariant for such systems). Further, by considering long trajectories of multidimensional systems, which start and end in some small region in phase space, and which are not quite periodic in general, we can formally show that, in a Poincaré cycle time or in the limit $T \rightarrow \infty$ the long-path action W is an adiabatic invariant for quasiperiodic and chaotic trajectories of multidimensional systems. Ehrenfest [95] has previously shown that W is an adiabatic invariant for multidimensional periodic systems. For ergodic systems, the phase volume $\Omega(E)$ is an adiabatic invariant (see next paragraph), and since the adiabatic invariant is apparently unique for such systems (Kasuga [90]), it should be possible to derive the adiabatic invariance of W from that of Ω for these systems, but we have

been so far unable to do this [86]. We note that counterparts of the Hannay angle have been established for adiabatically evolving nonintegrable [88] and ergodic [89] systems.

For one limiting case, complete chaos (ergodic motion) where the energy is the only constant of the motion, and long trajectories typically cover essentially the whole energy surface, a rigorous adiabatic invariant has been found [90, 91, 93], i.e. the phase volume $\Omega(E)$, enclosed by the energy surface $H(q, p) = E$:

$$\Omega(E) = \int dqdp \Theta[E - H(q, p)], \quad (5.31)$$

where Θ is the step function, which is unity for positive argument. (For multidimensional systems, $dqdp$ stands for $dq_1 dq_2 \dots dq_f dp_1 dp_2 \dots dp_f$.)

If $H(q, p, \lambda(t))$ depends on a slowly varying parameter $\lambda(t)$ so that $E = E(\lambda(t))$, then $\Omega(E(\lambda(t))) = \Omega(E(\lambda(0)))$. The adiabatic invariance of $\Omega(E)$ is intuitively plausible [94]. First consider the initial ($t = 0$) energy surface $H(q, p, \lambda(0)) = E(\lambda(0))$, and its image under an arbitrary time development (i.e. not necessarily slow). The final (image) surface at the time T is not in general a constant energy surface, and hence not in general a dynamical surface. However, by Liouville's theorem the volume enclosed by the final surface is equal to that enclosed by the initial surface. If the time-development from $t = 0$ to $t = T$ is sufficiently slow then at each stage the surface which has evolved from the $t = 0$ surface is a dynamical surface, by definition of adiabatic evolution, where allowed motions transform into allowed motions [95]. In this case the final $t = T$ surface is a true constant energy dynamical surface, and hence $\Omega(E(\lambda(T))) = \Omega(E(\lambda(0)))$ from Liouville's theorem.

For ergodic systems, there are no good quantum numbers except for the energy-ordering quantum number $N = 1, 2, \dots$, where $E_1 < E_2 < \dots$. A rough but useful semiclassical quantization scheme for ergodic systems based on $\Omega(E)$ is the Weyl rule [93]

$$\Omega(E_N) = N h^f, \quad (5.32)$$

which implicitly defines E_N . This is just a restatement of the early rule due to Planck [96] that each quantum state occupies a volume of size h^f in classical phase space. In practice, various improvements such as replacing N by $(N + \alpha)$ as in (2.4) are used [93]. More subtle improvements can be obtained from Gutzwiller's trace formula [28]. These relate fluctuations from

the smooth dependence of E on N predicted by (5.32) to the classical periodic orbits. In cases such as the x^2y^2 potential, where the classical $\Omega(E)$ diverges [98] due to the infinite channels of the potential (see Fig.1), one can estimate the smooth part of $N(E)$ (the cumulative number of states up to E) quantum mechanically [97] at low energy by an asymptotic expansion [98], with leading terms $\varepsilon^{3/2} \log \varepsilon$ and $\varepsilon^{3/2}$, where $\varepsilon = E(m^2/\hbar^4 C)^{1/3}$ is dimensionless. Application of the generalization of (5.32), i.e. $N(E_N) = N$, then gives reasonable estimates of the low-lying state energies [36, 98].

5.3 The Classical Hellmann - Feynman Theorem

Another consequence of the UMP (5.4) is the classical version of the Hellmann - Feynman theorem [99]

$$\frac{\partial E(I, \lambda)}{\partial \lambda} = \langle \frac{\partial H(q, p, \lambda)}{\partial \lambda} \rangle_I. \quad (5.33)$$

Here $H(q, p, \lambda)$ depends on a parameter λ , so that the energy $E(I, \lambda)$ depends on λ , as well as on the (constant) action I , and $\langle \dots \rangle_I$ denotes a time-average over a period $T(I)$ at fixed action I . We assume a one-dimensional periodic system for simplicity.

The quantum version of the theorem [100] is nowadays much better known, i.e.

$$\frac{\partial E_n(\lambda)}{\partial \lambda} = \langle \frac{\partial \hat{H}(\lambda)}{\partial \lambda} \rangle_n, \quad (5.34)$$

where $\langle \dots \rangle_n = \langle n | \dots | n \rangle$, and where the Hamiltonian operator $\hat{H}(\lambda)$ depends on some parameter λ , so that the energy eigenvalue $E_n(\lambda)$ and eigenfunction $\psi_n(\lambda)$ do also, because of

$$\hat{H}(\lambda)\psi_n(\lambda) = E_n(\lambda)\psi_n(\lambda). \quad (5.35)$$

Proof of (5.34) is simple; we apply $\partial/\partial\lambda$ to both sides of the relation $E_n(\lambda) = \langle \psi_n(\lambda) | \hat{H}(\lambda) | \psi_n(\lambda) \rangle$, and note that the two terms involving $\partial\psi_n(\lambda)/\partial\lambda$ add to zero because of (5.35), the hermiticity of $\hat{H}(\lambda)$, and the fact that we assume $\langle \psi_n(\lambda) | \psi_n(\lambda) \rangle = 1$ for all λ .

It is clear that (5.33) is the classical limit of (5.34), since as $\hbar \rightarrow 0$ (or $n \rightarrow \infty$) $\langle \dots \rangle_n \equiv \langle n | \dots | n \rangle$ approaches a time average over a period, and fixing n corresponds to fixing I , since $I \simeq n\hbar$. To derive (5.33) from (5.4) we consider $H(\lambda(t))$ to evolve adiabatically, so that λ and T in (5.4) can be

related. The derivation is intricate, however, and it is simpler to proceed instead from the RMP (2.3) for conservative systems. We apply (2.3) to two periodic trajectories of $H(\lambda)$, the true one with period $T(\lambda)$, and a virtual one taken as the true trajectory of $H(\lambda + \Delta\lambda)$ with period $T(\lambda + \Delta\lambda)$. Since first-order variations in $\bar{E} \equiv \langle H(\lambda) \rangle$ vanish (at fixed $W = 2\pi I$), we have

$$\langle H(\lambda) \rangle_{\lambda+\Delta\lambda} = \langle H(\lambda) \rangle_{\lambda} + O(\Delta\lambda^2), \quad (5.36)$$

where $\langle \dots \rangle_{\lambda}$ is an average over the trajectory with period $T(\lambda)$, and it is understood that I is the same for both $\langle \dots \rangle_{\lambda}$ and $\langle \dots \rangle_{\lambda+\Delta\lambda}$. From (5.36) we get for $\Delta\bar{E} \equiv \langle H(\lambda + \Delta\lambda) \rangle_{\lambda+\Delta\lambda} - \langle H(\lambda) \rangle_{\lambda}$ to $O(\Delta\lambda)$ accuracy

$$\Delta\bar{E} = \langle H(\lambda + \Delta\lambda) \rangle_{\lambda+\Delta\lambda} - \langle H(\lambda) \rangle_{\lambda+\Delta\lambda} = \langle H(\lambda + \Delta\lambda) - H(\lambda) \rangle_{\lambda+\Delta\lambda}.$$

Again neglecting $O(\Delta\lambda^2)$ terms we have

$$\Delta\bar{E} = \langle H(\lambda + \Delta\lambda) - H(\lambda) \rangle_{\lambda}. \quad (5.37)$$

Eq(5.37) yields $\partial\bar{E}/\partial\lambda = \langle \partial H(\lambda)/\partial\lambda \rangle_{\lambda}$, and since $\bar{E} = E$ for the true trajectory we get $\partial E/\partial\lambda = \langle \partial H(\lambda)/\partial\lambda \rangle_{\lambda}$, the Hellmann-Feynman relation (5.33) apart from notation.

As a simple example of (5.33) consider a 1D harmonic oscillator, with Hamiltonian $H(x, p) = p^2/2m + kx^2/2$ and $E(I) = I\omega$, where $\omega = (k/m)^{1/2}$ is the angular frequency. Choose $\lambda = k$, the force constant. The left side of (5.33) is $E/2k$ and the right side is $\langle x^2/2 \rangle = \langle V \rangle / k$. Thus we have $\langle V \rangle = E/2$, as is well known from the virial theorem [5]. If we choose $\lambda = m$, we find similarly $\langle K \rangle = E/2$.

As a second example, consider the bound motion of a 1D system with Hamiltonian $H(x, p) = p^2/2m + V(x)$. We change variables $x, p \rightarrow x', p'$ via the transformation $x' = \lambda x$, $p' = p/\lambda$, where λ is an arbitrary scale factor. This transformation is canonical and preserves the Hamiltonian [101], i.e. $H(x, p) = H_{\lambda}(x', p')$, where

$$H_{\lambda}(x', p') = \lambda^2 \frac{p'^2}{2m} + V\left(\frac{x'}{\lambda}\right). \quad (5.38)$$

The energy of a given state of motion is unchanged by the change from H to H_{λ} for any value of λ . We now use (5.38) in the Hellmann-Feynman relation (5.33). The left hand side vanishes. The right hand side derivative is

$$\frac{\partial H_{\lambda}}{\partial \lambda} = \frac{\lambda p'^2}{m} - \frac{x'}{\lambda^2} V' = \frac{2}{\lambda} K - \frac{x}{\lambda} V',$$

where $V' = \partial V(x)/\partial x$, and we have reverted to the original variables in the second form. Hence we get

$$\langle 2K \rangle = \langle xV' \rangle, \quad (5.39)$$

i.e. the virial theorem for an arbitrary potential $V(x)$.

6 Variational Principles in Relativistic Mechanics

Hamilton's Principle is widely used in Classical Relativistic Mechanics and in the Classical Theory of Electromagnetic and Gravitational Fields to derive covariant equations of motion (see, e.g.[102]). As for Maupertuis' Principle, it is widely believed that it is not well suited for that purpose because when operating with the energy one loses explicit covariance. In this section we demonstrate, first, how to use Maupertuis' Principle for the derivation of the covariant equations of motion and, the second, how to use it for the solution of specific problems.

Consider a relativistic particle with mass m and charge e moving in an arbitrary external electromagnetic field, described by a four-potential with contravariant components $A^\alpha = (A_0, A_i) = (\phi, A_i)$, and covariant components $A_\alpha = (A_0, -A_i) = (\phi, -A_i)$, where ϕ and A_i (for $i = 1, 2, 3$) are the usual scalar and vector potentials. Hamilton's Action for this system can be written in the Lorentz invariant form [102, 103]

$$S = m \int ds + e \int A_\alpha dx^\alpha, \quad (6.1)$$

where we use the sign of Lanczos [103] for S , opposite to that of Landau and Lifshitz [102]. The sign can be chosen arbitrarily; our choice allows us to use below the orthodox definitions $p = \partial L/\partial v$ and $H = pv - L$. The four-dimensional path runs from the initial point $x_A = (x_A^0, x_A^1, x_A^2, x_A^3)$ to the final point $x_B = (x_B^0, x_B^1, x_B^2, x_B^3)$ in four-dimensional space-time with corresponding proper times s_A and s_B . Here ds is the infinitesimal interval of the path (or of the proper time) $ds^2 = dx_\alpha dx^\alpha = g_{\alpha\beta} dx^\alpha dx^\beta = dx_0^2 - dx_i^2 = dt^2 - dx_i^2$, the metric has signature $(+, -, -, -)$ and we use the summation convention and units with $c = 1$, where c is the velocity of light. S is itself not gauge invariant, but a gauge transformation $A_\alpha \rightarrow A_\alpha + \partial f/\partial x^\alpha$ (for

arbitrary f) adds only constant boundary points terms to S , so that δS is unchanged. The HP is thus gauge invariant.

If we introduce a parameter τ along the four-dimensional path (proper time s along the true or any particular virtual path are valid choices) the action S can be rewritten in the form

$$S = m \int [v_\alpha v^\alpha]^{1/2} d\tau + e \int A_\alpha v^\alpha d\tau, \quad (6.2)$$

where $v^\alpha = dx^\alpha/d\tau$. In general, τ is both frame-independent and path-independent. A path-independent parameter is here invaluable for variational purposes, since we then do not have to vary the parameter when we vary the path. After the variations have been performed, we can then choose $\tau = s$, etc. The limits of the integrals are the invariants τ_A and τ_B . With respect to parameter τ we get the covariant Lagrangian

$$L = m[v_\alpha v^\alpha]^{1/2} + ev_\alpha A^\alpha, \quad (6.3)$$

and corresponding conjugate momenta

$$p^\alpha = \frac{\partial L}{\partial v_\alpha} = mv^\alpha/[v_\beta v^\beta]^{1/2} + eA^\alpha, \quad (6.4)$$

and Hamiltonian

$$H = p_\alpha v^\alpha - L = 0. \quad (6.5)$$

Thus in this particular covariant treatment the Hamiltonian is trivial. The dynamics is hidden in a constraint. One can see that the four conjugate momenta are not independent variables but satisfy the constraint

$$(p_\alpha - eA_\alpha)(p^\alpha - eA^\alpha) = m^2, \quad (6.6)$$

which is obvious from (6.4). This constraint can be taken into account with the help of the Lagrange multiplier method, i.e. with the help of an effective Hamiltonian [103, 108]

$$H_{eff} = \lambda[(p_\alpha - eA_\alpha)(p^\alpha - eA^\alpha) - m^2]. \quad (6.7)$$

We easily find that the Hamilton equations of motion with H_{eff} are equivalent to

$$\lambda = 1/2m$$

(for $\tau = s$, the true trajectory proper time) and to the covariant Lorentz equations of motion for a charged particle in an external field:

$$m \frac{dv_\alpha}{ds} = e F_{\alpha\beta} v^\beta,$$

$$F_{\alpha\beta} = \frac{\partial A_\beta}{\partial x^\alpha} - \frac{\partial A_\alpha}{\partial x^\beta}, \quad (6.8)$$

i.e. the same equations as are found from the Lagrangian (6.3) (see, e.g. [102]).

On the other hand the Hamilton equations with H_{eff} are equivalent to the MP or to the RMP with the same H_{eff} . Thus one can use any of the four Variational Principles (HP, RHP, GMP, RMP) either with Lagrangian (6.3) or with effective Hamiltonian (6.7) for the derivation of the covariant equations of motion. It is to be noted that in the covariant formulation of relativistic mechanics, the Hamiltonian (a Lorentz scalar) is not equal to the energy (the time component of a four-vector). Thus $\bar{E}$ in the Maupertuis Principle must be interpreted as

$$\bar{E} \equiv \bar{H}_{eff} = \frac{1}{\Delta\tau} \int_{\tau_A}^{\tau_B} d\tau H_{eff}, \quad (6.9)$$

where $\Delta\tau = \tau_B - \tau_A$. By contrast, in the noncovariant formulation discussed below, the Hamiltonian is equal to the energy $K + V + m$.

There are other, related, formal difficulties arising from (6.5). This condition ($H = 0$) is due to the fact that the Lagrangian L in (6.3) is homogeneous and of first degree in v_α (i.e. essentially linear in velocities v_α), so that $v_\alpha \partial L / \partial v_\alpha = L$ and hence the difference $v_\alpha \partial L / \partial v_\alpha - L$ (the Hamiltonian) vanishes [104]. Since $H = 0$, the Legendre transform relation (6.5) between L and H simplifies to

$$L = p_\alpha v^\alpha, \quad (6.10)$$

and hence the Legendre transform relation between W and S , i.e. $W - S = \bar{H} \Delta\tau$ (where $\bar{H} \equiv \bar{E}$), simplifies to

$$S = W = \int_{\tau_A}^{\tau_B} p_\alpha v^\alpha d\tau = \int_{x_A}^{x_B} p_\alpha dx^\alpha. \quad (6.11)$$

Because $H \equiv 0$, direct application of the GMP $(\delta W)_{\bar{E}=0} = 0$ would be cumbersome, and direct application of the RMP $(\delta \bar{E})_W = 0$ would be impossible.

All these difficulties can be traced to the well studied difficulty [5, 103, 105, 106, 107] of applying an action principle with kinematic [109] non-holonomic constraints, such as (6.6) for the momenta, or for the velocities $v_\alpha v^\alpha = 1$ for proper time s , or $v_\alpha v^\alpha = (ds/d\tau)^2$ for arbitrary parameter τ .

One way out of the difficulties is to introduce the constraint with a Lagrange multiplier for either the Lagrangian or Hamiltonian (as is done above for H_{eff}). Another way out is to exploit the nonuniqueness of the Lagrangian. For example [105], the covariant Lagrangian L (6.3) is equivalent to the covariant Lagrangian L' , where (with τ as parameter)

$$L' = \lambda(\tau) \frac{m}{2} v_\alpha v^\alpha + e v_\alpha A^\alpha, \quad (6.12)$$

where $\lambda(\tau) = d\tau/ds$. It is easily verified that (6.12) generates the same Euler-Lagrange equations of motion as (6.3). For $\tau = s$ (the proper time of the path in question), (6.12) simplifies to

$$L' = \frac{m}{2} v_\alpha v^\alpha + e v_\alpha A^\alpha, \quad (6.13)$$

which is often employed [5, 106]. The advantage of (6.12) and (6.13) is that they are nonhomogeneous in v_α , and hence will generate a nontrivial corresponding Hamiltonian H' . For (6.13), for example, we find

$$H' = \frac{1}{2m} (p_\alpha - e A_\alpha) (p^\alpha - e A^\alpha). \quad (6.14)$$

Note that H' (6.14) and H_{eff} (6.7) (for $\tau = s$) differ only by a constant. Use of the covariant L' and H' in place of L and H removes the difficulties of using all four variational principles (HP, RHP, GMP, RMP). For L and H , only the first two can be used. Note again that $\bar{E}$ is computed from H' as in (6.9), with H' in place of H_{eff} .

Now we come to the question of the solution of specific physical problems. In this case explicit covariance usually is not of great importance and it is more useful to find an appropriate convenient frame. For any given frame we can choose $\tau = t$ (the coordinate time) and obtain the complete set of nontrivial quantities

$$L = -m(1 - v^2)^{1/2} - e\phi + e\mathbf{v} \cdot \mathbf{A}, \quad S = \int_0^T L dt,$$

$$\mathbf{p} = \frac{m\mathbf{v}}{(1-v^2)^{1/2}} + e\mathbf{A},$$

$$W = \int_0^T \mathbf{p} \cdot \mathbf{v} dt = \int_0^T dt \left[\frac{mv^2}{(1-v^2)^{1/2}} + e\mathbf{v} \cdot \mathbf{A} \right], \quad (6.15)$$

$$H = \frac{m}{(1-v^2)^{1/2}} + e\phi, \quad \bar{E} = \frac{1}{T} \int_0^T H dt.$$

The expression for L in (6.15) is derived from (6.3) or (6.12) by choosing $\tau = t$, except that we have used the freedom to change the sign of L to agree with the standard choice in the noncovariant case [5]. Again we have chosen $t_A = 0$ and $t_B = T$. Note that the nonrelativistic form $W = \int_0^T 2K dt$ for the Maupertius action is not valid here, and also note that the Hamiltonian H is equal to the energy $K + e\phi + m$, despite the fact that $L \neq K - V$.

As a result the complete set of four variational principles is available in any particular frame. It is matter of taste or practical convenience to use one principle or another. From a practical point of view there is no difference in the usage of variational principles in nonrelativistic mechanics and in relativistic mechanics in a given frame.

As an example of the usage of the GMP for relativistic mechanics consider the motion of an electron in a uniform magnetic field $\mathbf{B} = (0, 0, B)$. A corresponding vector potential in the Coulomb gauge ($\nabla \cdot \mathbf{A} = 0$) is $\mathbf{A} = \mathbf{B} \times \mathbf{r}/2$, or $\mathbf{A} = (-yB/2, xB/2, 0)$ [116]. The solution of (6.8) in this case is well known - the electron moves with constant velocity along the direction of the field (along the z-axis in our case) and along a circular orbit in the xy plane with the cyclotron frequency

$$\omega = \frac{|e|B}{E}, \quad (6.16)$$

where E is the total energy of the particle including the kinetic energy of the motion in the z -direction.

To calculate this frequency from the GMP we take as a trial trajectory

$$x = A \cos \omega t, \quad y = A \sin \omega t, \quad v_z = 0,$$

where ω and A are free parameters. After some simple algebra we get

$$\bar{E} = \frac{m}{(1 - \omega^2 A^2)^{1/2}},$$

$$W = 2\pi A^2 \left[\frac{m\omega}{(1 - \omega^2 A^2)^{1/2}} - \frac{|e|}{2} B \right]. \quad (6.17)$$

In this case it seems that an economical method is to use the GMP (2.2). For this purpose one has to rewrite W as a function of $\bar{E}$

$$W = \frac{2\pi}{\omega^2} \left[1 - \left(\frac{m}{\bar{E}} \right)^2 \right] [\omega \bar{E} - \frac{|e|}{2} B], \quad (6.18)$$

and to calculate the variation δW at fixed $\bar{E}$. The GMP (2.2) translates here to $(\partial W / \partial \omega)_{\bar{E}} = 0$. One then easily finds the analytical solution

$$\omega = \frac{|e|B}{\bar{E}}.$$

The exact solution corresponds to the case when $\bar{E} = E$. For our trial trajectory, we happen to have $\bar{E} = E$ and the approximate solution coincides with the exact one (similar to the case of a harmonic oscillator).

This simple example demonstrates how to use variational principles for relativistic systems.

7 Classical Limit of Quantum Variational Principles

In section 2.1 we mentioned that the classical limit of the Schrödinger Quantum Variational Principle is the Reciprocal Maupertuis Principle for periodic or other steady-state motions, i.e.

$$\left(\delta \frac{\langle \psi | \hat{H} | \psi \rangle}{\langle \psi | \psi \rangle} \right)_n = 0 \xrightarrow{n \gg 1} \left(\delta \frac{1}{T} \int_0^T H(q, p) dt \right)_w = 0, \quad (7.1)$$

where $\hat{H}$ is the Hamiltonian operator corresponding to the classical Hamiltonian $H(q, p)$. Here T is the period for periodic motions, or $T \rightarrow \infty$ for other steady-state motions (quasiperiodic or chaotic). Solution of the left-hand side yields [118] the stationary state ψ_n with the energy eigenvalue E_n , i.e.

$$\hat{H}\psi_n = E_n\psi_n, \quad (7.2)$$

the time-independent Schrödinger equation. Details of the proof of (7.1) (which is straightforward only for integrable systems) are given in [12]. In ref.[119] it is explained how Schrödinger, in his first paper on wave mechanics in 1926, in essence worked backwards, i.e. starting from the right hand side of (7.1), he “derived” the left hand side, and thereby discovered wave mechanics. In his second paper on wave mechanics in 1926 he retracted this argument because his version of the right hand side of (7.1) was not quite correct. In this second paper he presented an alternative argument as a basis for wave mechanics (now in the standard textbooks) based on the analogy between geometric and wave optics on the one hand, and particle and wave mechanics on the other.

We consider now the generalization of (7.1) to nonstationary states $\psi(t)$, where the true dynamical states satisfy the time-dependent Schrödinger equation

$$i\hbar\partial_t\psi(t) = \hat{H}\psi(t), \quad (7.3)$$

where $\partial_t = \partial/\partial t$, and the initial state $\psi(0)$ is given. For notational simplicity we consider a single particle in three dimensions, so that $\psi = \psi(\mathbf{r}, t)$, and

$$\hat{H} = K + V = -\frac{\hbar^2}{2m}\nabla^2 + V(\mathbf{r}), \quad (7.4)$$

where $K = \mathbf{p}^2/2m$ and $\mathbf{p} = (\hbar/i)\nabla$. The arguments which follow are also valid for nonconservative systems, where $\hat{H} = \hat{H}(t)$.

We shall consider here the quantum time-dependent variational principle due to Frenkel and Dirac [120, 121, 122] for the wave function $\psi(t)$, but we note the existence of other quantum time-dependent variational principles, e.g. for the time-evolution operator and transition amplitudes [123], for the Heisenberg operator version [124] and for the matrix mechanics version [125] of the Hamilton Principle, for the density matrix [126], for the mean value of an arbitrary observable [127], and for time-dependent density functional theory [128]. (For completeness, we note that there are also other [130] quantum time-independent variational principles, in addition to that of Schrödinger.)

The original Frenkel-Dirac variational principle is [120]

$$\langle \delta\psi | \hat{H} - i\hbar\partial_t | \psi \rangle = 0, \quad (7.5)$$

for arbitrary variations $\delta\psi(\mathbf{r}, t)$ around the true dynamical state $\psi(\mathbf{r}, t)$. Since (7.5) is true for all possible variations $\delta\psi$, then $(\hat{H} - i\hbar\partial_t)\psi = 0$ for the true

states, as in (7.3). By adding (7.5) and its complex conjugate and using the hermiticity of $\hat{H}$ we can transform (7.5) to [129]

$$\delta \langle \psi | \hat{H} - i\hbar \partial_t | \psi \rangle = -i\hbar \langle \psi | \delta \psi \rangle . \quad (7.6)$$

Integrating over an arbitrary but fixed time interval $(0, T)$, and assuming $\psi(t=0)$ is fixed, we find [131] from (7.6)

$$\delta \int_0^T \langle \psi | \hat{H} - i\hbar \partial_t | \psi \rangle dt = -i\hbar \langle \psi(T) | \delta \psi(T) \rangle . \quad (7.7)$$

Note that (7.7) is not in the traditional form for a variational principle $\delta \int L dt = 0$, because of the right hand side. It is more like the unconstrained principles discussed earlier. Some authors drop the right hand side without comment [132], but it is required if arbitrary variations $\delta \psi$ in ψ are allowed; otherwise (7.7) will not generate the Schrödinger equation (7.3) and its complex conjugate [129]. It is possible to obtain a vanishing right-hand side by restricting the variations $\delta \psi$ in various ways [133], e.g. requiring the trial wave functions $\psi(t)$ to be normalized at all times. Another way is to write the right hand side as $-i\hbar \langle \delta \psi(T) | \psi(T) \rangle^*$, formally regard ψ and its complex conjugate ψ^* as independent (see below), and assume $\psi^*(T)$ (as well as $\psi(0)$) is fixed. However, we find it simplest to retain the general form (7.7), with no constraints on $\psi(t)$ except for the fixing of the initial condition $\psi(0)$. (When dealing explicitly with bound states later, we assume boundary conditions on $\psi(\mathbf{r}, t)$.)

By writing out the scalar products $\langle A | B \rangle \equiv \int d\mathbf{r} A(\mathbf{r})^* B(\mathbf{r})$ in (7.7) and using the hermiticity of $\mathbf{p} = (\hbar/i)\nabla$, we can write (7.7) in field theory form

$$\delta \int_0^T dt \int d\mathbf{r} \mathcal{L} = -i\hbar \int d\mathbf{r} \psi(T)^* \delta \psi(T) , \quad (7.8)$$

where the Lagrangian density $\mathcal{L}$ is

$$\mathcal{L}(\psi, \psi^*) = \frac{\hbar^2}{2m} |\nabla \psi|^2 + V|\psi|^2 - i\hbar \psi^* \partial_t \psi . \quad (7.9)$$

Since ψ is a complex function, whose real and imaginary parts can be varied independently, it is well known [134] that ψ and ψ^* can be regarded as independent for variational purposes. These two variations in (7.8) with $\mathcal{L}$ given by (7.9) will generate the Schrödinger equation and its complex conjugate.

Following Feynman [132], we change variables $\psi, \psi^* \rightarrow A, S$ (or ρ, S) in the Lagrangian density $\mathcal{L}(\psi, \psi^*)$, where $A(\mathbf{r}, t)$ is the amplitude and $S(\mathbf{r}, t)$ the phase of the wave function, via the Madelung transformation [135]:

$$\psi = Ae^{iS/\hbar} = \sqrt{\rho}e^{iS/\hbar} , \quad (7.10)$$

where $\rho = A^2 = |\psi|^2$ is the probability density. In terms of ρ and S we have

$$\mathcal{L}(\rho, S) = \rho \frac{(\nabla S)^2}{2m} + \frac{\hbar^2}{2m} (\nabla \sqrt{\rho})^2 + \rho V + \rho \partial_t S - \frac{i\hbar}{2} \partial_t \rho. \quad (7.11)$$

In terms of ρ, S the boundary term in (7.8) is

$$\langle \psi(T) | \delta \psi(T) \rangle = \int d\mathbf{r} \sqrt{\rho(T)} \delta \sqrt{\rho(T)} + \frac{i}{\hbar} \int d\mathbf{r} \rho(T) \delta S(T) . \quad (7.12)$$

We now take the classical ($\hbar \rightarrow 0$) limit of (7.8). Using (7.11) and (7.12) we get

$$\delta \int_0^T dt \int d\mathbf{r} \rho \left[\frac{(\nabla S)^2}{2m} + V + \partial_t S \right] = \int d\mathbf{r} \rho(T) \delta S(T) . \quad (7.13)$$

We have assumed that in the limit $\hbar \rightarrow 0$, ρ and S approach well defined limits (which we continue to denote by ρ and S). In particular we assume that $\nabla \sqrt{\rho}$ does not contain an $O(\hbar^{-1})$ term [136]. We vary ρ and S independently in (7.13). Varying ρ , with S fixed, gives

$$\int \int dt d\mathbf{r} \delta \rho \left[\frac{(\nabla S)^2}{2m} + V + \partial_t S \right] = 0.$$

Since $\delta \rho(\mathbf{r}, t)$ is arbitrary, the bracket [...] must vanish, so that

$$\partial_t S + \frac{(\nabla S)^2}{2m} + V = 0 . \quad (7.14)$$

This is the classical Hamilton-Jacobi equation [138] for the classical action $S(\mathbf{r}, t) = \int_{\mathbf{r}_0, 0}^{\mathbf{r}, t} L dt'$ along true paths.

Next we vary S in (7.13) with ρ fixed. We integrate the $\delta(\nabla S)^2$ term by parts using

$$\int d\mathbf{r} \rho (\nabla \delta S) \cdot \nabla S = - \int d\mathbf{r} \delta S \nabla \cdot (\rho \nabla S) , \quad (7.15)$$

where we drop a surface term by assuming $\rho \rightarrow 0$ for $\mathbf{r} \rightarrow \infty$. This assumes the corresponding quantum state has finite norm. We integrate the $\delta(\rho\partial_t S)$ term by parts using

$$\int_0^T dt \rho \partial_t \delta S = \rho(T) \delta S(T) - \int_0^T dt \delta S \partial_t \rho , \quad (7.16)$$

where we use $\delta S(t=0) = 0$ which follows from the fixing of $\psi(t=0)$. Using (7.15) and (7.16) in (7.13) thus gives

$$\int_0^T dt \int d\mathbf{r} [\nabla \cdot (\rho \frac{\nabla S}{m}) + \partial_t \rho] \delta S = 0.$$

Since this is true for arbitrary $\delta S(\mathbf{r}, t)$, we get

$$\partial_t \rho + \nabla \cdot (\rho \frac{\nabla S}{m}) = 0 . \quad (7.17)$$

This is the continuity equation, reflecting conservation of total probability, since [138] $\nabla S = \mathbf{p} = m\mathbf{v}$ is the momentum of the particle. Note that if the right hand side of (7.13) were missing, we would not get the correct continuity equation. In contrast to what we have done here, Feynman [132] keeps the $\hbar$ terms in (7.11), and the variational principle then leads to quantum generalizations [137] of the Hamilton - Jacobi (7.14) and continuity (7.17) equations.

Thus we see that the classical limit of the quantum time-dependent variational principle is the classical variational principle for the Hamilton-Jacobi and continuity equations. Interestingly the classical limit of the quantum time-dependent variational principle leads to a classical variational principle for the Hamilton action S along the true trajectories, whereas the classical limit of the quantum time-independent variational principle in the form of the left side of (7.1) leads to a classical variational principle for the trajectories themselves (RMP -the right side of (7.1)). An alternative form [139] of the quantum time-independent variational principle,

$$\delta \langle \psi | \hat{H} - E | \psi \rangle = 0, \quad (7.18)$$

will generate in the classical limit a variational principle for the Maupertuis (or time-independent) action $W(\mathbf{r})$ along the true paths. In (7.18) E is a constant Lagrange multiplier introduced to remove the normalization constraint $\langle \psi | \psi \rangle = 1$. The extremizing $\psi(\mathbf{r})$'s which satisfy (7.18) obey

$$(\hat{H} - E)\psi = 0, \quad (7.19)$$

so that E is the energy eigenvalue corresponding to the eigenfunction ψ .

To derive the classical limit of (7.18) we proceed much as before. (We assume $\hat{H}$ is time-independent in this section.) We first apply the Madelung transformation to $\psi(\mathbf{r})$, i.e.

$$\psi(\mathbf{r}) = A(\mathbf{r})e^{iW(\mathbf{r})/\hbar} , \quad (7.20)$$

where $A(\mathbf{r})$ is the amplitude and $W(\mathbf{r})$ the phase. Following similar steps as above, we find (7.18) becomes in the classical ($\hbar \rightarrow 0$) limit

$$\delta \int d\mathbf{r} \rho(\mathbf{r}) \left[\frac{(\nabla W)^2}{2m} + V - E \right] = 0 , \quad (7.21)$$

where $\rho(\mathbf{r}) = A(\mathbf{r})^2 = |\psi(\mathbf{r})|^2$.

Varying $\rho(\mathbf{r})$ in (7.21) with $W(\mathbf{r})$ fixed gives

$$\frac{(\nabla W)^2}{2m} + V - E = 0 , \quad (7.22)$$

the time-independent Hamilton-Jacobi equation [138] for the Maupertuis action $W(\mathbf{r}) = \int_{\mathbf{r}_0}^{\mathbf{r}} \mathbf{p} \cdot d\mathbf{r}'$ along true paths. If we vary $W(\mathbf{r})$ in (7.21) with fixed $\rho(\mathbf{r})$ we get the continuity equation

$$\nabla \cdot \left(\rho \frac{\nabla W}{m} \right) = 0 , \quad (7.23)$$

since [138] $\nabla W = \mathbf{p} = m\mathbf{v}$ is the momentum of the particle.

In ref.[12], we show how the classical limit of (7.18) can be taken in a different way (using standard 1D WKB wave functions) and for a 1D periodic orbit leads to $\delta(S + ET) = 0$, i.e. the UHP for a periodic orbit of period T .

We note that, just as for the right hand side of (7.1), the derivation of the classical limits (7.14) and (7.22) is somewhat formal for the chaotic motions of nonintegrable systems, since classical values of S and W may not exist as well defined functions for such systems. Unlike the nonrelativistic limit ($v/c \rightarrow 0$), which is regular, the limit $\hbar/S \rightarrow 0$ or $\hbar/W \rightarrow 0$ is singular, which makes the classical limit subtle and interesting [140].

8 Summary and Conclusions

We summarize the two groups of classical variational principles reviewed in this paper. First we have the Maupertuis principles:

$$\delta W - T\delta\bar{E} = 0, \quad (\delta W)_{\bar{E}} = 0, \quad (\delta\bar{E})_W = 0. \quad (8.1)$$

The first of these is the unconstrained form (UMP), the second the general Maupertuis principle (GMP), and the third is the reciprocal Maupertuis principle (RMP). By Legendre transformation of the principles in (8.1) we obtain the Hamilton principles:

$$\delta S + E\delta T = 0, \quad (\delta S)_T = 0, \quad (\delta T)_S = 0. \quad (8.2)$$

The first is the unconstrained form (UHP), the second the Hamilton principle (HP), and the third the reciprocal Hamilton principle (RHP). In all these principles, the end-positions of the trajectory q_A and q_B are held fixed, and the notation indicates what else is fixed and what is varied.

We have given a number of examples showing how to use these principles to solve practical problems, focussing particularly on the three Maupertuis principles. Both classical and semiclassical applications have been given, and some comparisons are made with the results from quantum variational principle calculations. The classical limits of two quantum variational principles (time-dependent and time-independent) are given. In particular, the RMP is the classical limit of the Schrödinger time-independent quantum variational principle. The RMP is thereby naturally adapted to semiclassical applications

In the applications, the variational principles (both classical and quantum) are solved by the direct variational method, where one guesses a trial solution containing adjustable parameters, and adjusts the parameters to satisfy the variational principle. The Euler-Lagrange differential equation is not required in this method. When we choose a simple trial solution, the results can be found simply and analytically.

Using the direct variational method the RMP and UMP have been used to estimate semiclassically the energy levels in several examples. The levels $E_{n_1, n_2}, \dots$ can be expressed in analytical form, and the accuracy is quite acceptable in most cases, even for nonintegrable systems which are usually treated by much more complicated semiclassical methods, e.g. the Gutzwiller trace formula [28], or quantizing the Birkhoff-Gustavson normal form [141]. It would be useful to have comparisons of the results from the Maupertuis Principle-based method with those from the latter two and other methods for a number of specific problems. The methods should be compared for accuracy of results and computational complexity. Some limited comparisons of the Maupertuis Principle and normal form methods have been done for the x^2y^2 oscillator of sec. 4.1 in refs. [58, 68], and limited comparisons for

this model of results from the Maupertuis Principle and the semiclassical adiabatic methods [35] are presented in Table 1 of sec. 4.1.1, but much more extensive comparisons would be of interest.

Variational principles appeal to various individuals for differing reasons, both aesthetic [142] and practical. We have stressed the utility for classical and semiclassical particle dynamics calculations in this review. Engineers too make heavy use of variational principles in solving practical problems of classical continuum mechanics [143]. On rare occasions variational principles have even led to new laws of physics [144].

9 Acknowledgements

We gratefully acknowledge the Natural Sciences and Engineering Research Council of Canada for financial support of this work. We thank Donald Sprung for a careful reading of the manuscript, David Garfinkle for helpful remarks on relativistic mechanics and Stepan Bulanov for help with TeX.

10 Appendix I. Variational Notation and Reciprocal Principles

“... since δy is the variation of y with x kept constant, we could conveniently write it as $(\delta y)_x$, in accordance with a convention used in thermodynamics... It is curious that in spite of the obvious need in partial differentiation for precise statement of what is being kept constant, such statement is not embodied in customary notation of pure mathematics, though it is provided in thermodynamics,...

H. Jeffreys and B.S. Jeffreys
Methods of Mathematical Physics
Cambridge U.P., 1946.

Variational problems ask for the function which makes stationary an integral involving the function and its derivatives. Usually there is a constraint which is fixed. For example, we are to find a function $y(x)$, describing a closed planar curve, of fixed length L , which maximizes the area A enclosed

by the curve $y(x)$. A and L are functionals of $y(x)$: $A = A[y(x)]$, $L = L[y(x)]$. For our purpose here, we do not need the precise form of A or L . L is the constraint and A is the functional to be varied. We use the following notation to describe this variation

$$(\delta A)_L = 0, \quad (\text{A1.1})$$

where the constraint L is indicated as a subscript on the bracket enclosing the variation δA of the functional A . The variation δA is first-order in δy , and is called the first variation when higher-order variations are considered [15]. The constraint L is denoted explicitly in (A1.1) because it is the most important, or the one we wish to focus on. There may be other constraints which are left implicit, e.g. that the planar curve pass through a given point (x_B, y_B) . In this paper we always leave fixed end-position constraints on curves (closed or open) implicit.

The solution of the problem (A1.1) is the function $y(x)$, which satisfies a differential equation, the Euler-Lagrange equation. The reciprocal variational problem has the constraint L and the varied functional A interchanged. In other words, it is described by the equation

$$(\delta L)_A = 0. \quad (\text{A1.2})$$

The two equations, (A1.1) and (A1.2), have the same solution $y(x)$. The reciprocal variation is searching for the function $y(x)$ which makes stationary (here minimizes) the length L of a closed curve with fixed enclosed area A . The common solution for both problems is a circle, as is clear intuitively. One way to prove reciprocity is to use a trick, due to Lagrange, called the method of Lagrange multipliers. The problem (A1.1) is first restated in unconstrained form as

$$\delta A - \lambda \delta L = 0, \quad (\text{A1.3})$$

where the variations of both A and L are unconstrained and λ is an undetermined multiplier. To obtain (A1.2) from (A1.3), we simply specify that we now fix A (i.e. set $\delta A = 0$) and vary L , i.e. $(\delta L)_A = 0$. This proves the reciprocity theorem.

Reciprocity is clear intuitively in the isoperimetric problem (A1.1). An example of reciprocity from physics is the Gibbs conditions for equilibrium in thermodynamics [117]. The reciprocal conditions $(\delta S)_U = 0$ and $(\delta U)_S = 0$ are both conditions for thermal equilibrium, where S is the entropy and U the internal energy. These express the facts that the state of thermal equilibrium

is that state which maximizes the entropy for fixed internal energy, and also is that state which minimizes the internal energy for fixed entropy. (The unconstrained form [145] is $\delta U - T\delta S = 0$, where the Lagrange multiplier is the temperature T . Here the implicit constraints in all cases are fixed system volume V and mole numbers N_α) Reciprocal VP's in mechanics have only recently been studied [12, 13] - see Sec.2.

This is all we shall need from the calculus of variations; for more details one should consult a text-book [146].

11 Appendix II. Equivalence of the Maupertuis and General Maupertuis Principles

In Sec. 2.2, for conservative systems, we discussed the equivalence of the Hamilton Principle (HP)

$$(\delta S)_T = 0, \quad (\text{A2.1})$$

with the General Maupertuis Principle (GMP)

$$(\delta W)_{\bar{E}} = 0. \quad (\text{A2.2})$$

Here we show directly the equivalence of the GMP (A2.2) with the original Maupertuis Principle (MP)

$$(\delta W)_E = 0. \quad (\text{A2.3})$$

In all these principles the end-positions q_A and q_B are held fixed. Since (A2.1), (A2.2) and (A2.3) all generate the same (Euler-Lagrange) equations of motion, it is clear they must be equivalent, and it is interesting to show directly the equivalence of (A2.2) and (A2.3). We must show that relaxing the constraint of fixed E in (A2.3) to fixed $\bar{E}$ in (A2.2) does not generate spurious or unphysical solutions.

The constraint of fixed $E = H(q, p)$ in (A2.3) applies along and between the trial trajectories. To relax this constraint, we use the method of Lagrange multipliers [147]. Since $H(q, p) = \text{const}$, we have $\delta H = 0$, and hence

$$\lambda \int_0^T \delta H dt = 0,$$

where λ is an arbitrary constant. The trajectories all start at $t = 0$ at fixed point A and end at $t = T$ (which may vary with the trial trajectory) at fixed

point B . We release the constraint of fixed E in (A2.3) and choose λ such that

$$\delta W = \lambda \int_0^T \delta H dt, \quad (\text{A2.4})$$

where δ indicates a variation from the true trajectory value and $\delta H \neq 0$ in (A2.4). This can be rewritten as

$$\delta W = \lambda [\delta \int_0^T H dt - H(T) \delta T], \quad (\text{A2.5})$$

where $H(T) \equiv H(q(T), p(T))$, and where we recall that the end-time T is not fixed in (A2.3). Recalling the definition $\bar{E} = \int_0^T H dt / T$ enables us to write (A2.5) as

$$\delta W = \lambda \delta(T \bar{E}) - \lambda E \delta T, \quad (\text{A2.6})$$

where we have used the fact that $H = E = \text{const}$ on the true trajectory. Expanding (A2.6) gives

$$\delta W = \lambda \bar{E} \delta T + \lambda T \delta \bar{E} - \lambda E \delta T. \quad (\text{A2.7})$$

The first and third terms on the right hand side of (A2.7) cancel, since $\bar{E} = E = \text{const}$ on the true trajectory, as assumed in the MP (A2.3). This means

$$\delta W = \lambda T \delta \bar{E}. \quad (\text{A2.8})$$

To see that $\lambda = 1$ in (A2.8), we apply (A2.8) to the special case of two true trajectories, with actions W and $W + dW$ and energies E and $E + dE$. For this case (A2.8) reads $\partial W / \partial E = \lambda T$. Comparison of this with the standard relation [5] $\partial W / \partial E = T$ gives $\lambda = 1$. Hence we have

$$\delta W = T \delta \bar{E}, \quad (\text{A2.9})$$

which is the unconstrained Maupertuis Principle (UMP) (2.10). Fixing $\bar{E}$ in (A2.9) then gives (A2.2), the GMP.

Thus (A2.2) follows from (A2.3). The converse, the derivation of (A2.3) from (A2.2) is easy; we simply restrict the trial trajectories in (A2.2) to those of fixed energy E , since the equations of motion following from (A2.2) imply energy conservation. Hence (A2.2) and (A2.3) are equivalent.

We note once more that conservation of energy is assumed in the original MP (A2.3), whereas it is a consequence of the general MP (A2.2).

References

- [1] Nesbet R K 2003 *Variational Principles and Methods In Theoretical Physics and Chemistry* (New York: Cambridge U.P.); L.R. Ram-Mohan L R 2002 *Finite Element and Boundary Element Applications in Quantum Mechanics* (Oxford: Oxford U.P.); Drake G W F 1998 *Encyclopedia of Applied Physics* (New York: Wiley-VCH) **23** 121; Gerjuoy E, Rau A R P and Spruch L 1983 *Rev.Mod.Phys.* **55** 725; Epstein S T 1974 *The Variation Method in Quantum Chemistry* (New York: Academic).

Quantum VP's for scattering states are reviewed in Adhikari S K 1998 *Variational Principles for the Numerical Solution of Scattering Problems* (New York: Wiley). Classical VP's for scattering are discussed in ref. [12].

- [2] For example, in Yourgrau and Mandelstam [9] the notation is reversed. Hamilton himself used the current notation, but, for the actions along true paths, termed S and W the principal and characteristic functions, respectively.
- [3] There is a recent exception: Papastavridis J G 2002 *Analytical Mechanics* (New York: Oxford U.P.), p. 1044, where the general Maupertuis principle is discussed. This book contains a masterly and historical account of the various variational principles.
- [4] Arnold V I 1989 *Mathematical Methods of Classical Mechanics* 2nd edition (New York: Springer Verlag)
- [5] Goldstein H, Poole C and Saffko I 2002 *Classical Mechanics* 3rd edition (New York: Addison Wesley)
- [6] Sommerfeld A 1964 *Mechanics* translated from the 4th German edition (New York and London: Academic Press)
- [7] In this review we focus on global (integral) VP's. We do not discuss the various local (differential) VP's due to d'Alembert, Gauss, Hertz, Gibbs, Jourdain and others [3, 9, 103]. Jourdain's principle, and its connections to others, is reviewed by Wang L-S and Pao Y-H 2003 *Am. J. Phys* **71** 72

- [8] For example, Ité U E 2001 *Global Thinking and Local Action* (Aldershot, Hampshire: Ashgate Publishing)
- [9] Yourgrau W and Mandelstam S 1968 *Variational Principles in Dynamics and Quantum Theory* 3rd edition (London: Pitman, Philadelphia: W.B.Saunders), reprinted, Dover, New York, 1979; Terral M 2002 *The Man Who Flattened the Earth* (Chicago: University Of Chicago Press)
- [10] quoted in footnote 77 of Arnold, ref. [4], p 246
- [11] C.G.J.Jacobi 1884 *Vorlesungen über Dynamik*, zweite ausgabe, herausgegeben von A.Clebsch (Braunschweig/ Wiesbaden: F.Vieweg), reprinted by Chelsea 1969, p 44. The quotation in this text is more explicit than [10]: it names the textbooks of Poisson, Lagrange and Laplace as some of the best!
- [12] Gray C G, Karl G and Novikov V A 1996 *Ann. Phys.* **251** 1
- [13] Gray C G, Karl G and Novikov V A 1996 *Am. J. Phys.* **64** 1177
- [14] Percival I C 1977 *Adv. Chem. Phys.* **36** 1, and references therein
- [15] van Brunt B 2004 *The Calculus of Variations* (New York: Springer) Ch 10
- [16] Karl G and Novikov V A 1995 *Phys. Rev. D* **51** 5069; 1995 *J. Exp. Theor. Phys.* **80** 783
- [17] Holonomic (coordinate) constraints are assumed to be taken into account by the choice of the $q_1, q_2, \dots, q_f$. Nonholonomic (velocity) constraints (see note [109]) are assumed absent in the early sections but are considered briefly in sec.6 and App.II.
- [18] We also exclude quasi-coordinates (or nonholonomic coordinates), e.g. for a rigid body, the three rotation angles around space-fixed xyz axes. VP's using such variables are discussed by Papastavridis [3], and by Marsden J E and Ratiu T S 1994 *Introduction to Mechanics and Symmetry* (New York: Springer) p 390. A seminal paper is Poincaré H 1901 *C.R.Acad.Sci.* **132** 369

- [19] The condition $H = K + V$ is valid for normal systems, for which $L = K - V$, the holonomic constraints (if any) are time-independent and the reference frame is inertial. If these conditions are satisfied K is quadratic in the $\dot{q}$'s. Note that the single condition K quadratic in the $\dot{q}$'s is not sufficient to have $H = K + V$ since for some systems with time-dependent constraints K is quadratic in the $\dot{q}$'s: see, e.g. de Castro A S 2000 *Eur. J. Phys.* **21** 23 ; Ferrario C and Passerini A 2002 *Eur. J. Phys.* **22** 11.
- When $H \neq K + V$, the generalized Maupertuis principles require $\bar{H}$ in place of $\bar{E}$, where $H = p\dot{q} - L$ (see Secs. 5.2.1 and 6).
- [20] Percival I C 1974 *J. Phys. A* **7** 794 ; ibid 1979 **12** 157 ; see also Klein A and Li C-t 1979 *J. Math. Phys.* **20** 572. Percival later extended his VP for invariant tori to cantori: 1980 *Amer. Inst. Phys. Conf. Proc.* **55** 302. A cantorus is, roughly speaking, a torus-shaped surface with holes such that what remains forms a Cantor set of points. A trajectory confined to a torus surface is quasiperiodic. A trajectory starting on a cantorus will slowly “leak away” because of the holes, hence is not confined to the surface so that the quasiperiodicity is destroyed in the long run, and is thus weakly chaotic. See Arnold et al [108], p.194, for a precise definition. Also see Percival I C 1991 in *Chaos and Quantum Physics* eds. Giannoni M-J, Voros A and Zinn-Justin J 1991 (Amsterdam: North-Holland) p 11
- [21] Greenberg W R, Klein A and Li C-t 1995 *Phys. Rev. Lett.* **75** 1244
- [22] Goldstein et al, ref. [5], p 54; Low F 1997 *Classical Field Theory* (New York: Wiley) p 282
- [23] Goldstein et al, ref. [5], p 597; Lanczos, ref. [103], p 402; Oliver D 1994 *The Shaggy Steed of Physics* (New York: Springer) p 39; Schwinger J, De Raad L L Jr., Milton K A and Tsai W-Y 1998 *Classical Electrodynamics* (Reading MA: Perseus Books) p 90
- [24] See, for example, Feynman R P 1948 *Rev. Mod. Phys.* **20** 267, and references therein.
- [25] The unconstrained Hamilton Principle (2.8) is closely related to what Hamilton termed the law of varying action (see Landau and Lifshitz [83],

p.134), and also to what some authors refer to as the Weiss action principle: see Sudarshan E C G and Mukunda N 1974 *Classical Dynamics* (New York: Wiley), reprinted by Krieger 1983, p 16.

If end-position variations are allowed, as well as end-time variations, the UHP (2.8) generalizes to $\delta S = (p_B \delta q_B - H_B \delta t_B) - (p_A \delta q_A - H_A \delta t_A)$, where p_B is the true value of p at end-point B , etc, and for conservative systems $H_A = H_B = E$. (This generalization can be derived from the general first variational theorem [29] of variational calculus, which includes end-point variations - see Appendix A of [12] for a related discussion of the UMP.) What Hamilton termed the law of varying action is the special case for the variation between two true trajectories, so that $\delta S = dS$, $\delta p_i = dp_i$ and $\delta t_i = dt_i$ are exact differentials.

- [26] This particular analogy can be pushed further. Just as there are numerous other thermodynamic potentials besides the Helmholtz free energy F and the Gibbs free energy $G = F + pV$, (where p is the pressure and V the volume), e.g. the internal energy $U = F + TS$, (where T is the temperature and S the entropy), we can introduce in mechanics further actions via further Legendre transformations. For example, define $S' = S - p_B q_B$, where q_B is the final end-point position and p_B the true momentum at B . If we relax the constraint of fixed q_B , the UHP generalizes further to $\delta S = -E \delta T + p_B \delta q_B$. In terms of S' this transforms to $\delta S' = -E \delta T - q_B \delta p_B$. In terms of S' , the usual fully constrained HP is $(\delta S')_{T, p_B} = 0$, replacing $(\delta S)_{T, q_B} = 0$. Here, as always, the constraint on q_A is still left implicit in both. Further Legendre (using p_A, q_A and/or W) and reciprocal transformations are obvious. In this review we focus on the four basic variational principles summarized at the end of sec.2.2.
- [27] Szamosi G 1973 *Found. Phys.* **3** 241; Peterson M A 1979 *Am. J. Phys.* **47** 488; Toffoli T in 1999 *Feynman and Computation* ed. Hey A J G (Reading MA: Perseus Books) p 349; Bailyn M 1990 *A Survey of Thermodynamics* (New York: AIP Press) p 416. In particular, authors interested in linking the principle of least action to the second law of thermodynamics include (see Bailyn) Boltzmann, Clausius, Helmholtz, Ehrenfest [95] and Planck: see also, e.g., Planck M 1915 *Eight Lectures on Theoretical Physics* (New York: Columbia Univ. Press), reprinted by Dover, 1998, p 97; Thomson J J 1888 *Applications of Dynamics to Physics and Chemistry* (London: Macmillan) reprinted by Dawson, 1968; Fényes I

1952 *Zeit. f. Phys.* **132** 140, and de Broglie L 1995 *Diverses Questions de Mécanique et de Thermodynamique Classiques et Relativistes* (Berlin: Springer)

- [28] Gutzwiller M C 1990 *Chaos in Classical and Quantum Mechanics* (New York: Springer); Child M S 1991 *Semiclassical Mechanics with Molecular Applications* (Oxford: Oxford U.P.); Brack M and Bhaduri R K 1997 *Semiclassical Physics* (Reading MA: Addison-Wesley)

- [29] Gelfand I M and Fomin S J 1963 *Calculus of Variations* (Englewood Cliffs, New Jersey: Prentice Hall) Ch 8

- [30] The direct method described in the text originates with Ritz (precursors due to Hamilton and Rayleigh).

Euler first described the method of discretizing time in the action integral, treating the $q(t_i)$ as discrete variables, and obtaining discretized (finite difference) versions of the Euler-Lagrange (and Newton) equations, to be solved with the two end-positions and the duration fixed. This method has also been much used in recent years: see, e.g., Percival 1991 [20]; Beck T L, Doll J D and Freeman D L 1989 *J. Chem. Phys.* **90** 3181; Basille A G and Gray C G 1992 *J.Comp.Phys.* **101** 80; Marsden J E and West M 2001 *Acta Numerica* **10** 357; Monforte J C 2002 *Geometric, Control and Numerical Aspects of Nonholonomic Systems* (New York: Springer) p 143; Elber R, Cárdenas A, Ghosh A and Stern H A 2003 *Adv. Chem. Phys.* **126** 93.

Courant R (Bull. Amer. Math. Soc. **49**, 1 (1943)) was one of the first to suggest spacial discretization of variational functionals involving fields $\phi(\mathbf{r})$, a precursor to the modern finite element method (Ram-Mohan, ref.[1]) which uses a basis of localized trial functions.

- [31] Unfortunately, in some good textbooks the authors misapply the original MP $(\delta W)_E = 0$ for 1D problems by employing trial trajectories which do not conserve the energy E : see, e.g., Tabarrok and Rimrott [115], p. 180, or Park D 1990 *Classical Dynamics and its Quantum Analogues* 2nd ed.(New York: Springer) p 80.

- [32] McWeeny R and Coulson C A 1948 *Proc. Camb. Phil. Soc.* **44** 413

- [33] Crane H R 1995 *Am. J. Phys.* **63** 33 .

- [34] Olsson M G 1978 *Am. J. Phys.* **46** 1118; *ibid* 1981 **49** 531; Synge J L and Griffiths B A 1959 *Principles of Mechanics* 3rd ed. (New York: McGraw Hill) p 335; Routh E J 1898 *A Treatise on the Dynamics of a Particle* (Cambridge: Cambridge U.P.), reprinted by Dover 1960 p 349; Airy G B 1851 *Phil.Mag.* Series 4 **2** 141.
- In the second edition of *Mécanique Analytique* (1815), Lagrange gave an incorrect result for eq (3.11). For an English translation of the second edition see Lagrange J L 1997 *Analytical Mechanics*, eds. Boissonnade A and Vagliente V N (Dordrecht: Kluwer) p 462. The error was corrected by Bertrand, the editor of the third edition (1855), following a communiqué by Bravais.
- [35] Martens C C, Waterland R L, and Reinhardt W P 1989 *J.Chem.Phys.* **90** 2328
- [36] Whelan N D 1997 *J.Phys.A* **30** 533
- [37] Dahlqvist P and Russberg G 1990 *Phys. Rev. Lett.* **65** 2837
- [38] The RMP method yields $\omega_x/\omega_y = W_y/W_x$ and hence after semiclassical quantization $\omega_x/\omega_y = (n_y + \frac{1}{2})/(n_x + \frac{1}{2})$, the same as (4.7) found by the fully quantum VP. Since ω_x/ω_y is therefore rational, the optimum quantized trajectories are periodic, at least at this level of approximation. Periodic trajectories also play a key role in Gutzwiller's trace formula method of semiclassical quantization [28].
- [39] Tinkham M 1964 *Group Theory and Quantum Mechanics* (New York: McGraw Hill) p 325
- [40] Rajan M and Junkins J L 1983 *Int. J. Nonlinear Mech.* **18** 335
- [41] Bukta D, Karl G and Nickel B G 2000 *Can. J. Phys.* **78** 449
- [42] Szeredi T and Goodings D A 1992 *Phys. Rev. Lett.* **69** 1640; *ibid.* 1993 *Phys. Rev.E* **48** 3529
- [43] Burnell F, Mann R B and Ohta T 2003 *Phys. Rev. Lett.* **90** 134101
- [44] Pauli W 1973 *Pauli Lectures in Physics* vol.5 (Cambridge: MIT Press)

- [45] Shinohara H 2000 *Rep. Prog. Phys.* **63** 843; Eletsii A V 2000 *Uspekhi* **43**, 111
- [46] Dresselhaus M S, Dresselhaus G and Eklund P S 1996 *Science of Fullerenes and Carbon Nanotubes* (San Diego: Academic Press)
- [47] Cioslowski J 1995 *Electronic Structure Calculations on Fullerenes and Their Derivatives* (New York: Oxford U.P.)
- [48] Joslin C G, Gray C G, Poll J D, Goldman S and Buckingham A D 1995 *Collision and Interaction-Induced Spectroscopy* eds Tabisz G C and Neuman M N NATO ASI Ser. C **452** (Dordrecht: Kluwer) p 261
- [49] Joslin C G, Yang J, Gray C G, Poll J D and Goldman S 1995 *Lectures on Thermodynamics and Statistical Mechanics* eds Castas M, Rodriguez R and Benavides A L (Singapore: World Scientific) p 146
- [50] Exohedral complexes XC_n , where X is on the outside of the cage, have also been studied, e.g. H_2C_{60} [51] and H_2C_∞ (nanotubes) [52]. Complexes such as $K_3 C_{60}$ are superconductors in the solid state [46].
- [51] FitzGerald S A, Forth S and Rinkoski M 2002 *Phys. Rev. B* **65** 140 302(R)
- [52] Williams K A, Pradhan B K, Eklund P C, Kostov M K and Cole M W 2002 *Phys. Rev. Lett.* **88** 165 502; Kostov M K, Cheng H, Herman R M, Cole M W and Lewis J C 2002 *J. Chem. Phys.* **116** 1720
- [53] Joslin C G, Gray C G, Goddard J D, Goldman S, Yang J and Poll J D 1993 *Chem. Phys. Lett.* **213** 377; Joslin C G, Yang J, Gray C G, Goldman S, and Poll J D 1993 *Chem. Phys. Lett.* **208** 86; Joslin C G, Yang J, Gray C G, Goldman S, and Poll J D 1993 *Chem. Phys. Lett.* **211** 587; Joslin C G, Gray C G, Goldman S, Yang J and Poll J D 1993 *Chem. Phys. Lett.* **215** 44
- [54] Hernández-Rojas J, Bretón J and Gomez Llorente J M 1997 *J. Phys. Chem. Solids* **58** 1689, and references therein
- [55] Tellgmann R, Krawez N, Lin S-H, Hertel I V and Campbell E E B 1996 *Nature* **382** 407

- [56] Gromov A, Ostrovskii D, Lassensson A, Jönsson M and Campbell E E B 2003 *J.Phys.Chem.B* **107** 11290
- [57] Joslin C G, Gray C G and Goldman S 1995 *Chem. Phys.Lett.* **244** 93; ibid 1996 **251** 414 (E)
- [58] Lee-Dadswell G R and Gray C G 1995 *Can. J. Phys.* **78** 599
- [59] Lee-Dadswell G R and Gray C G 2003 *Can. J. Phys.* submitted
- [60] Mandziuk M and Bacic Z 1994 *J. Chem. Phys.* **101** 2126
- [61] Joslin C G, Gray C G and Goldman S 1994 *Chem. Phys.Lett.* **227** 405
- [62] Cioslowski J, Rao N, Pernal K and Moncrieff D 2003 *J. Chem. Phys.* **118** 4456
- [63] Melle-Franco M, Kuzmany H and Zerbetto F 2003 *J. Phys.Chem.B* **107** 6986 and references therein.
- [64] For $\text{Li}^+@\text{C}_{60}$ and $\text{LiH}@\text{C}_{60}$, anisotropic perturbations of the isotropic potential, and the consequent chaotic motions, have been studied by Hernández-Rojas et al. [54].
- [65] Yakobson B I and Smalley R E 1997 *American Scientist* **85** 324
- [66] Bug A L R, Wilson A and Voth G A 1992 *J. Phys. Chem.* **96** 7864; Cardini G, Procacci P, Salvi P R and Schettino V 1992 *Chem. Phys. Lett.* **200** 39; Li Y S and Tománek D 1994 *Chem. Phys. Lett.* **221**, 453
- [67] The potential $V(r)$ can be fit [48] to simple analytic forms (polynomial, Morse, etc). In some cases (e.g. $\text{Li}^+@\text{C}_{60}$) the potential has its minimum off-centre. For $\text{K}^+@\text{C}_{60}$ the potential is essentially $V(r) = Cr^4$, a rare example of a pure quartic oscillator in nature.
- [68] Lee-Dadswell G R 1999 *M.Sc.thesis* University of Guelph, p 58
- [69] Leubner C and P. Krumm P 1990 *Eur. J. Phys.* **11** 31 ; Anderson N and Arthurs A M 1997 *Eur. J. Phys.* **18** 404 ; Kobe et al. [70]; del Valle G, Campos I and Jiménez J L 1996 *Eur. J. Phys.* **17** 253

- [70] Bateman H 1931 *Phys. Rev.* **38** 815; Havas P 1957 *Nuovo Cimento Suppl.* **5** 363; Leitmann G 1963 *J. Appl. Mech.* **30** 623; Denman H H 1966 *Am. J. Phys.* **34** 1147; Ray J R 1974 *Am. J. Phys.* **47** 626; Greenberger D M 1979 *Int. J. Math. Phys.* **20** 762; Kobussen J 1979 *Acta Phys. Aust.* **51** 293; Bahar L Y and Kwatny H G 1981 *Am. J. Phys.* **49** 1062; Dekker H 1981 *Phys. Rep.* **80**, 1; Lemos N A 1981 *Am. J. Phys.* **49** 1181; Kobe D H, Reali G and Sieniutycz S 1986 *Am. J. Phys.* **54** 997; Huang Y-S and Lin C-L 2002 *Am. J. Phys.* **70** 741; Vujanovic B D and Jones S E 1989 *Variational Methods in Nonconservative Systems* (Boston: Academic Press); Edelen D G B 1977 *Lagrangian Mechanics of Nonconservative Nonholonomic Systems* (Leyden: Noordhoff); Soper D E 1976 *Classical Field Theory* (New York: Wiley) p 126; Santilli R M 1978 *Foundations of Theoretical Mechanics* Vol. 1 (New York: Springer) pp. 183, 199, 206; Logan J D 1977 *Invariant Variational Principles* (New York: Academic Press) p 56.

These workers go beyond the Rayleigh dissipation function method [5], and incorporate the friction into the Lagrangian (or Hamiltonian). For example, for a harmonic oscillator with linear damping (equation of motion $m\ddot{x} + \beta\dot{x} + kx = 0$), one easily verifies that a suitable Lagrangian is $L(x, \dot{x}, t) = \exp(\beta t/m)[m\dot{x}^2/2 - kx^2/2]$, and the corresponding Hamiltonian is $H(x, p, t) = \exp(-\beta t/m)p^2/2m + \exp(\beta t/m)kx^2/2$.

- [71] Buch L H and Denman H H 1976 *Phys. Lett. A* **55** 325. See also Yourgrau and Mandelstam [9], p.47, who note essentially (5.3) without remarking explicitly that it is valid for nonconservative systems. Similarly see Landau and Lifshitz [83], p.139, Routh [34], p.301, and Boltzmann [72], p. 121, where the nonconservative aspect is more explicit.
- [72] Boltzmann L 1904 *Vorlesungen über die Principe der Mechanik II Teil* (Leipzig: Barth), reprinted by Wissenschaftliche Buchgesellschaft, Darmstadt 1974, along with Part I (1897).
- [73] The form $\delta W = \int_0^T \delta H dt$ for the UMP, which is easily shown to be equivalent to (5.4), essentially appears in passing in the equations of Yourgrau and Mandelstam [9], p.46, but they do not note explicitly its validity for nonconservative systems. For the special case of periodic systems evolving adiabatically with parameter $\lambda(t)$, Ehrenfest [95] gives a relation related to (5.4); he has δE rather than $\delta \bar{E}$ (as in the original

MP), and our δT is replaced by a $d\lambda$ term. Ehrenfest attributes his result to Boltzmann and Clausius.

As before, in the UMP (5.4) we have fixed the initial time $t_A = 0$ and allow the final time $t_B \equiv T$ to vary. If both t_A and t_B are allowed to vary, (5.4) generalizes to $T\delta\bar{E} - \delta W - [E(t_B) - \bar{E}]\delta t_B + [E(t_A) - \bar{E}]\delta t_A = 0$. This reduces to (5.4) for fixed t_A , and to (2.10) for conservative systems ($E(t_A) = E(t_B) = \bar{E}$). Further generalizations, allowing q_B and/or q_A to vary, are discussed in notes [25, 26]

- [74] If we start the cycle at the moving wall, for example, instead of the fixed one, the action I_n is not an exact invariant. See Crawford F S 1990 *Am.J.Phys* **58** 337; Gignout C and Brut F 1989 *Am.J.Phys.* **57**, 422; Percival I and Richards D 1982 *Introduction to Dynamics* (Cambridge: Cambridge U.P.) p 142; Lochak and Meunier [90], p.18; Brack and Bhaduri [28], p. 10.
- [75] Whittaker E T 1937 *A Treatise on the Analytical Dynamics of Particles and Rigid Bodies* 4th edition (Cambridge: Cambridge U.P.), reprinted by Dover, 1944, p 248.
- [76] Lichtenberg A J and Lieberman M A 1992 *Regular and Chaotic Dynamics* 2 ed (New York: Springer) p 57 and 216. Krainov V P 2002 *Selected Mathematical Methods in Theoretical Physics* (London: Taylor and Francis) p 143.
- [77] Hannay J H 1985 *J.Phys.A* **18** 221; Berry M V 1985 *J.Phys.A* **18** 15. These papers and others on the Berry phase are reprinted in Shapere A and Wilczek F eds 1989 *Geometric Phases in Physics* (Singapore: World Scientific)
- [78] Berry M V and Morgan M A 1996 *Nonlinearity* **9** 787
- [79] Berry M V 1990 *Physics Today* Dec., p 34; Berry M V 1988 *Scientific American* Dec., p. 46.
- [80] Shapere A and Wilczek F, ref. [30], p 3; Marsden J E and Ratiu T S, ref. [18], p 44; Khein A and Nelson D 1992 *Am. J. Phys.* **61** 170; Kugler M 1989 *Am. J. Phys.* **57** 247; Montgomery R 1988 *Commun. Math. Phys.* **120** 269; Greenberger D M 1979 *J. Math. Phys.* **20** 762; Koiller J 1989 *Contemp. Math.* **97** 159

- [81] Grimsehl E 1932 *A Textbook of Physics* Vol. 1 (London: Blackie and Son) p 165; Sommerville W B 1972 *Quart. J. Roy. Astron. Soc.* **13** 40; Hart J B, Miller R E and Mills R L 1987 *Am. J. Phys.* **55** 67
- [82] Fowles G R and Cassiday G L 1999 *Analytical Mechanics* 6 ed.(Fort Worth: Saunders) p 196
- [83] Landau L D and Lifshitz E M 1969 *Mechanics* 2 ed. (Oxford: Pergamon) p 127
- [84] Child, ref. [29], p 75
- [85] Golin S 1988 *J. Phys. A* **21** 4535
- [86] The converse can be made plausible: from an assumed adiabatic invariance of the long path W , the adiabatic invariance of Ω follows. We use [87], for $T \rightarrow \infty$, $W \rightarrow \oint_C \sum_i p_i dq_i = \sum_i \int \int_{R_i} dq_i dp_i$, where R_i is the projected area of curve C on the q_i, p_i plane, and the fact that the dynamical invariance (Poincaré invariant) of $\Omega = \int \dots \int dq_1 dp_1 \dots dq_f dp_f$ can be derived [87] from the dynamical invariance of $\sum_i \int \int dq_i dp_i$,

The adiabatic invariance of long-path W 's for ergodic systems follows heuristically from the UMP (5.4) as described in the text. Alternatively, we can heuristically argue that for $T \rightarrow \infty$, the long path is essentially closed in phase space, so that the Poincaré dynamical invariant $W \equiv \oint \sum_i p_i dq_i$ remains essentially adiabatically constant as the energy surface and the path on it slowly evolve. Obviously the limiting procedures are delicate and problematic. Even if our conjecture is true, the adiabatic invariance may appear to be of the seemingly uninformative type $\infty = \infty$, since $W \rightarrow \infty$ for $T \rightarrow \infty$. To extract finite results, T will need to be taken as, say, the Poincaré cycle time. We note that, in another situation where $W \rightarrow \infty$ as $T \rightarrow \infty$, i.e. scattering orbits, useful results can still be extracted [12].

The arguments in the above paragraph, as well as the one in the text, apply also more generally to the chaotic motions of nonergodic nonintegrable systems. Since adiabatic invariants have not been found previously for such systems, it would be of interest to prove, or disprove, these suggestions. If proven, it would then be of interest to establish a general semiclassical quantization scheme for nonintegrable systems based on W (generalizing (5.32) which holds for ergodic systems).

- [87] Tabor M 1989 *Chaos and Integrability in Nonlinear Dynamics* (New York: Wiley) p 55 and p 86
- [88] Montgomery [45]; Golin S, Knauf A and Marmi S 1989 *Commun. Math. Phys.* **123** 95
- [89] Robbins J M and Berry M V 1992 *Proc. Roy. Soc. A* **436** 634
- [90] Hertz P 1910 *Ann.der Phys.* **33** 225, 537. Hertz's proof of the adiabatic invariance of the phase volume $\Omega(E)$ made explicit what was implicit in the statistical mechanical works of Boltzmann, Gibbs, and others. (Recall the Boltzmann relation for the entropy $S = k_B \log \Omega(E)$; reversible adiabatic processes ($dS = 0$) correspond to constant $\Omega(E)$.) Hertz's argument is available in a few textbooks: Kahan T 1960 *Physique Théoretique* Tome Premier, Vol. II, (Paris: Presses Universitaires) p 482; Becker R 1969 *Theory of Heat* (New York: Springer) p 131; Münster A 1969 *Statistical Thermodynamics*, Vol.I (Berlin: Springer) p 52; Toda M, Kubo R and Saito N 1983 *Statistical Physics I* (Berlin: Springer) p 48. For a rigorous proof, see Kasuga T 1961 *Proc. Japan Acad.* **37** 366, 372, 377: see also Lochak P and C.Meunier C 1988 *Multiphase Averaging for Classical Systems* (New York: Springer) p 239.
- [91] Reinhardt W P 1994 *Prog. Theoret.Phys. Suppl.* **116** 179, and references therein
- [92] Skodje R T and Cary J R 1988 *Computer Phys. Rep.* **8** 221
- [93] Berry M V, in 1985 *Theoretical and Applied Mechanics* eds. F.I.Niordson and N.Olhoff (Amsterdam: Elsevier) p 83, and references therein
- [94] Brown R, Ott E and Grebogi C 1987 *Phys. Rev. Lett.* **59** 1173
- [95] Ehrenfest P 1917 *Phil.Mag.* **33** 500. Reprinted in *Sources of Quantum Mechanics* ed. van der Waerden B L 1967 (Amsterdam: North Holland), reprinted by Dover 1968, p 79. An extended version of this paper is available in *Paul Ehrenfest Collected Scientific Papers*, ed. Klein M J 1959 (Amsterdam: North Holland) p 378.

Later Ehrenfest's student J.M.Burgers showed that the one-cycle (or torus) actions I_i are adiabatic invariants for multidimensional quasiperiodic systems (Burgers J M 1918 *Proc.Amsterdam Acad.* **20** 163; for a

textbook treatment, see Landau and Lifshitz [83], p 161.) The subtle conditions (concerning the nondegeneracy of the system frequencies ω_i during the adiabatic evolution) required for the I_i to be adiabatic invariants are discussed by Dirac P A M 1925 *Proc. Roy. Soc.* **107** 725 and von Laue M 1925 *Ann. der Phys.* **76** 619

- [96] Planck M 1915 *Verhand. der Deutsch. Physikal. Gesell.* **17** 407, 438
- [97] The cumulative number of states up to E , $N(E)$, is defined quantum mechanically by $N(E) = \sum_n \Theta(E - E_n) = \text{tr } \Theta(E - \hat{H})$, where $\hat{H}$ is the Hamiltonian operator and tr denotes a trace.
- [98] Tomsovic S 1991 *J.Phys. A* **24** 733
- [99] McRae S M and Vrscaj E R 1992 *J. Math. Phys.* **33** 3004; McKinley W A 1971 *Am. J. Phys.* **39** 405; Weisskopf V 1934 *Zeit.f.Phys.* **89** 27. For periodic systems Ehrenfest [95] showed that the classical Hellmann-Feynman theorem is equivalent to the adiabatic invariance of the action W . Ehrenfest (who was Boltzmann's student) attributes this result to Boltzmann and Clausius.

The classical theorem can also be derived (Epstein [100]) from classical perturbation theory, using the result (Ter Haar [111], p.156) that the first-order energy shift ΔE due to a small change in the Hamiltonian $H(\lambda)$ by the amount $(\partial H/\partial \lambda)\Delta \lambda$ is $\Delta E = \langle (\partial H/\partial \lambda)\Delta \lambda \rangle_\lambda$.

Our derivation, and that of Epstein [100], allow λ to occur anywhere in the Hamiltonian. Other explicit published derivations assume λ occurs in the potential. Implicit derivations are contained in a number of places: Van Vleck J H 1926 *Quantum Principles and Line Spectra*, *Bull. Nat.Res. Council*, Vol. 10 Part 4, p 205; Becker [90], p 129; Münster [90], p51; Landau and Lifshitz [83], p 155.

The classical and quantum dynamical Hellmann-Feynman relations can be generalized to statistical mechanical relations; for systems in thermal equilibrium we find $\partial F/\partial \lambda = \langle \partial H/\partial \lambda \rangle$, where F is the Helmholtz free energy and $\langle \dots \rangle$ denotes a (classical or quantum) canonical ensemble average. See Gubbins K E and Gray C G 1984 *Theory of Molecular Fluids* (Oxford: Oxford U.P.) p 236, and references therein. Thus in statistical mechanics we do not have $\partial \langle H \rangle/\partial \lambda = \langle \partial H/\partial \lambda \rangle$ in the canonical ensemble; the correct relation is $\partial \langle H \rangle/\partial \lambda = \langle \partial H/\partial \lambda \rangle$

$-\beta[< H\partial H/\partial\lambda > - < H > < \partial H/\partial\lambda >]$, where $\beta = 1/k_B T$, with k_B Boltzmann's constant and T the temperature. (In the microcanonical ensemble, where $H = < H > = E$ because of the absence of energy fluctuations, we do find $\partial E/\partial\lambda = < \partial H/\partial\lambda >$, both classically and quantum mechanically.)

- [100] Hellmann H 1937 *Einführung in die Quantenchemie* (Vienna: Denticke); Feynman R P 1939 *Phys. Rev.* **56** 340; see also Van Vleck J H 1928 *Phys. Rev.* **31** 587, and Pauli W 1933 *Handbuch der Physik* **24**, Part 1, eds. Geiger H and Scheel K (Berlin: Springer), translated edition 1980 *General Principles of Quantum Mechanics* (Berlin: Springer) p 86. For a review of other derivations, applications and extensions, see Epstein S T 1981, in *The Force Concept in Chemistry* (New York: Van Nostrand Reinhold) ed. Deb B M, p 1.

The Hellmann - Feynman theorem seems to be a good example of the Arnold - Berry “theorem”: any named result was actually found by somebody else.

- [101] Goldstein et al., ref. [5], p 371
- [102] Landau L D and Lifshitz E M 1962 *The Classical Theory of Fields* 2nd edition (New York: Pergamon) p 49
- [103] Lanczos C 1970 *The Variational Principles of Mechanics* 4th edition (Toronto: University of Toronto Press) reprinted by Dover Publications, New York, 1986, p 324.
- [104] The condition $H = 0$ also occurs nonrelativistically if a parameter τ is used in place of t ; see, e.g., Goldstein et al [5], p.320, or Lanczos [103], p. 186.
- [105] Doughty N A 1990 *Lagrangian Interaction* (Reading MA: Addison-Wesley) p 416; Leubner C 1986 *Eur. J. Phys.* **7** 17; Johns O D 1985 *Am. J. Phys.* **53** 982
- [106] Barut A O 1964 *Electrodynamics and Classical Theory of Fields and Particles* (London: Macmillan), reprinted by Dover, 1990, p 60.
- [107] Forbes G W 1991 *Am. J. Phys.* **59** 1130

- [108] Dirac P A M 1964 *Lectures on Quantum Mechanics* (New York: Yeshiva University and Academic Press) p 9; Arnold V I, Kozlov V V and Neishtadt A I 1988 in *Dynamical Systems III* ed. Arnold V I (Berlin: Springer) p 39
- [109] Nonholonomic constraints $f_i(q, \dot{q}, t) = 0$ contain the velocities $\dot{q}_\alpha$, such that the constraints cannot be integrated up to global coordinate constraints $g_i(q, t) = 0$. Kinematic nonholonomic constraints arise from introducing redundant variables as in sec.6 on relativistic mechanics, or by introducing constants of the motion as constraints [110] as in App.II for the MP, or by imposing the kinematic constraint between $\dot{q}$ and the velocity or the momentum after first having released the constraint [111]. Dynamic nonholonomic constraints arise from physical forces of constraint, such as arise in rolling motion [112]. Kinematic nonholonomic constraints can be handled by an effective Lagrangian $L_{eff} = L + \sum_i \lambda_i f_i$, where λ_i are Lagrange multipliers [103, 106, 107]. Dynamic nonholonomic constraints cannot be handled by such an effective Lagrangian (contrary to what is stated in some excellent textbooks) and construction of an HP or MP is highly problematic [115].
- [110] Kapuscik E and Uzes C A 1982 *Am. J. Phys.* **50** 1094; Pars L A 1962 *Introduction to Calculus of Variations* (London: Heineman) p 23
- [111] Leubner C and Marte M A M 1985 *Eur.J.Phys.* **6** 22; Whatley M C 1990 *Am.J.Phys.* **58** 1006; Ter Haar D 1964 *Elements of Hamiltonian Mechanics* 2nd ed. (Amsterdam: North Holland) p 100
- [112] Two classic rolling motion examples, both of which are integrable systems, are the nonslipping tippe top [113] and the nonslipping Euler disc (or spinning coin) [114].
- [113] Gray C G and Nickel B G 2000 *Am. J. Phys.* **68** 821, and references therein
- [114] Petrie D, Hunt J L and Gray C G 2002 *Am. J. Phys.* **70** 1025, and references therein
- [115] Papastavridis [3] gives a clear discussion and references to the historical literature; Tabarrok B and Rimrott F P J 1994 *Variational Methods and Complementary Formulation in Dynamics* (Dordrecht: Kluwer) p 142;

- Rund H 1966 *The Hamilton-Jacobi Theory in Calculus of Variations* (New York: Van Nostrand Reinhold), reprinted by Krieger 1973, p 352; Pars L A 1965 *A Treatise on Analytic Dynamics* (Cambridge: Cambridge U.P.), reprinted by Ox Bow Press, 1979, p 528; Rumiantsev V V 1978 *J. Appl. Math. and Mech.* **42**, 407; *ibid.* 1979 **43** 625; Whittaker [75], p 249; Saletan E J and Cromer A H 1970 *Am. J. Phys.* **38** 892
- [116] Another common choice is $\mathbf{A} = (0, xB, 0)$. This choice is known as the Landau gauge (in this gauge $\mathbf{A}$ is invariant under translations along the y axis). The first choice is known as the symmetric gauge (because of the rotational invariance of $\mathbf{A}$ about the z axis).
- [117] Gibbs J W 1906 *The Scientific Papers of J. Willard Gibbs, Vol.1, Thermodynamics* (London: Longmans, Green), reprinted by Dover, 1961, p 56. For modern treatments, see Callen H B 1985 *Thermodynamics* 2nd edition (New York: Wiley) p 133, and Bailyn [27], p 224.
- [118] There are various ways to constrain the variation so that ψ_n is selected. In principle, one can minimize the energy $\langle \psi | \hat{H} | \psi \rangle / \langle \psi | \psi \rangle$ with the trial ψ 's restricted to be orthogonal to the exact $\psi_0, \psi_1, \dots, \psi_{n-1}$. In practice, since $\psi_0, \psi_1, \dots, \psi_{n-1}$ are rarely known exactly, one can find an approximate ψ_n by choosing trial ψ 's with n nodes (for 1D problems). Alternatively, one can choose a trial ψ which is known to be the n th eigenstate of a similar Hamiltonian (see sec. 3.1 for an example).
- [119] Gray C G, Karl G and Novikov V A 1999 *Am. J. Phys.* **67** 959
- [120] Frenkel J 1934 *Wave Mechanics, Advanced General Theory* (Oxford: Oxford U.P.) p 253. Frenkel states that he took his argument from the appendix to the Russian edition of Dirac P A M 1930 *The Principles of Quantum Mechanics* (Oxford: Oxford U.P.), but we are unable to locate the argument in this reference. See also Heisenberg W 1930 *The Physical Principles of the Quantum Theory* (Chicago: University of Chicago Press), reprinted by Dover, 1960, p 163
- [121] A minimum residual principle

$$\delta ||(\hat{H} - i\hbar\partial_t)\psi||^2 \equiv \delta \langle (\hat{H} - i\hbar\partial_t)\psi | (\hat{H} - i\hbar\partial_t)\psi \rangle = 0$$

is similar to that of Frenkel and Dirac, and was proposed by McLachlan A D and Bell M A 1964 *Rev. Mod. Phys.* **36** 844.

- [122] For recent applications of the Frenkel-Dirac Variational Principle to molecular and condensed matter problems, see e.g. refs [133] and also Beck M H, Jackle A, Worth G A and Meyer H-D 2000 *Phys. Rep.* **324** 1; Wang H 2000 *J. Chem. Phys.* **113** 9948.
- [123] Lippmann B A and Schwinger J 1950 *Phys. Rev.* **79** 469; Gerjuoy et al. [1].
- [124] Schwinger J 1951 *Phys. Rev.* **82** 914. The generalized operator UHP is $\delta\hat{S} = \hat{p}_B\delta q_B - \hat{H}_B\delta t_B$ (allowing both final end parameters (q_B, t_B) to vary, and fixing the initial ones (q_A, t_A)), where $\hat{p}_B = \hat{p}(t_B)$, $\hat{H}_B = \hat{H}(t_B)$, $\hat{F}(t)$ is a Heisenberg time-dependent operator, $\hat{S} = \int_{t_A}^{t_B} \hat{L}dt$ the action operator, and $\hat{L}(\hat{q}(t), \dot{\hat{q}}(t))$ the Lagrangian operator. If $\delta q_B = 0$ we get the operator UHP, $\delta\hat{S} = -\hat{H}_B\delta t_B$, and if the end parameter variations both vanish, we get the operator HP, $\delta\hat{S} = 0$. From the generalized operator UHP Schwinger derives what is nowadays known as the Schwinger Action Principle for the variation of the propagator $\langle q_B, t_B | q_A, t_A \rangle \equiv \langle q_B, t_A | \hat{U}(t_B, t_A) | q_A, t_A \rangle$, where $\hat{U}(t_B, t_A)$ is the evolution operator, i.e. $\delta \langle q_B, t_B | q_A, t_A \rangle = (i/\hbar) \langle q_B, t_B | \delta\hat{S} | q_A, t_A \rangle$. Here an arbitrary number of intermediate state parameters $(q_i, t_i), (q'_i, t'_i), \dots$ are allowed to vary on the left hand side, e.g. q_i, t_i in $\langle q_B, t_B | q_A, t_A \rangle = \int dq_i \langle q_B, t_B | q_i, t_i \rangle \langle q_i, t_i | q_A, t_A \rangle$, as well as the final end-state parameters q_B, t_B . (Schwinger also discusses other possible variations). If there are no end-state parameter variations, then $\delta\hat{S} = 0$ and hence $\delta \langle q_B, t_B | q_A, t_A \rangle = 0$. The analogy of the last relation with the classical HP is clearer when we write the classical HP as $\delta S(q_B, t_B; q_A, t_A) = 0$.
- Schwinger's original paper discusses the case of relativistic quantum field theory. For a nonrelativistic quantum mechanics discussion, with examples, see Schwinger's posthumously published lecture notes, Schwinger J 2001 *Quantum Mechanics* ed Englert B-G (Berlin: Springer) p 207. For the connection to Feynman's path integral expression for the propagator, see Yourgrau and Mandelstam, ref. [9], p 139.
- [125] Greenberg et al. [21]; Born M and Jordan P 1925 *Z. Phys.* **34** 858, reprinted and translated into English in van der Waerden [95], p 289
- [126] Eboli O, Jackiw R and Pi S-Y 1988 *Phys. Rev. D* **37** 3557; Rajagopal A K 1997 *Phys. Lett.A* **228** 66

- [127] Balian R and Veneroni 1985 *Ann. Phys.* **164** 334
- [128] Runge E and Gross E K U 1984 *Phys. Rev. Lett.* **52** 997
- [129] Langhoff P W, Epstein S T and Karplus M 1972 *Rev. Mod. Phys.* **44** 602
- [130] For VP's for diagonal and off-diagonal matrix elements of arbitrary operators, and for VP's for wave-functions, see Gerjuoy et al [1] and references therein. For a ground state energy VP based on path integrals, see Feynman R P and Hibbs A R 1965 *Quantum Mechanics and Path Integrals* (New York: McGraw-Hill) p 303. For a VP for the ground state energy as a functional of the particle density (density functional theory), see Hohenberg P and Kohn W 1964 *Phys. Rev.* **136B** 864.
- [131] Titov A V 1993 *Int. J. Quant. Chem.* **45** 71
- [132] Feynman R P 1972 *Statistical Mechanics* (Reading MA: Benjamin) p 303
- [133] Löwdin P O and Mukherjee P K 1972 *Chem. Phys. Lett.* **14** 1; Kramer P and Saraceno M 1981 *Geometry of the Time-dependent Variational Principle in Quantum Mechanics* (New York: Springer); Deumens E, Diz A, Longo R and Öhrn Y 1994 *Rev. Mod. Phys.* **66** 917
- [134] Epstein, ref [1], p 20
- [135] Madelung E 1926 *Zeit. Phys.* **40** 322; de Broglie L 1926 *Compt. Rend.* **183** 447; Bohm D 1952 *Phys. Rev.* **85** 166
- [136] Holland P R 1993 *The Quantum Theory of Motion* (Cambridge: Cambridge U.P.)p 220; Rosen N 1964 *Am. J. Phys.* **32** 597
- [137] These are often referred to as the de Broglie-Bohm or quantum hydrodynamic equations: see, e.g., Holland [136], p 15 and p 69; Wyatt R E 2002 *J. Chem. Phys.* **117** 956; Kendrick B K 2003 *J. Chem. Phys.* **119** 5805, and references therein.

The quantum continuity equation retains the same form (7.17), and the quantum Hamilton-Jacobi equation takes the form (7.14) with an additional so-called quantum potential term $V_Q = -(\hbar^2/2m)\rho^{-1/2}\nabla^2\rho^{1/2}$ added to V . These equations can also be (and are usually) derived by

substituting the Madelung form (7.10) into the Schrödinger equation (7.3) and using (7.4).

- [138] Goldstein et al., ref. [5], Ch. 10
- [139] In practice (7.18) is less useful for bound states than (7.1) since E and ψ are both unknown for bound states. For scattering states E is known, and $\langle \psi | \psi \rangle \rightarrow \infty$ causes no problem in (7.18), as it does in (7.1).
- [140] Berry M V 2002 *Physics Today* May p 10; Batterman R W 2002 *The Devil in the Details: Asymptotic Reasoning in Explanation and Emergence* (Oxford: Oxford U.P.) p 109
- [141] See, e.g., Swimm R T and Delos J B 1979 *J.Chem.Phys.* **71** 1706; Jaffe C and Reinhardt W P 1982 *J.Chem.Phys.* **77** 5191.

In the present context, the normal form is a perturbation series, often generated by Poisson bracket and Lie transform methods, which expresses the system energy E as a power series in the good actions I_i , $E = \sum_i a_i I_i + \sum_{ij} a_{ij} I_i I_j + \dots$ Semiclassical quantization is achieved by the EBK substitution $I_i \rightarrow (n_i + \alpha_i)\hbar$.

In molecular physics the quantized version of the normal form is usually referred to as the Dunham expansion: see, e.g., Toda M 2002 *Adv. Chem. Phys.* **123** 153; Herman M, Lievin J, Auwera J V and Campargue A 1999, *Adv. Chem. Phys.* **108**, 1
- [142] Hildebrandt S and Tromba A 1996 *The Parsimonious Universe: Shape and Form in the Natural World* (New York: Springer)
- [143] Reddy J N 2002 *Energy Principles and Variational Methods in Applied Mechanics* (New York: Wiley)
- [144] The derivation of the Einstein gravitational field equations from the Einstein - Hilbert action is the classic example: see, e.g., Landau and Lifshitz [102], p 307
- [145] This is usually written as $\delta F = 0$, where $F = U - TS$ is the Helmholtz free energy: i.e. the free energy is a minimum at equilibrium at a given T , V and N_α .

- [146] For example, see Gelfand and Fomin [29], van Brunt [15], or Akhiezer H I 1988 *The Calculus of Variations* (London: Harwood)
- [147] Routh uses this method to, in effect, transform the original MP (A2.3) into the unconstrained Hamilton Principle (2.8), and from this he obtains the Euler-Lagrange equations of motions. Pars uses a similar method to derive Euler-Lagrange equations from the original MP; he however, allows λ to be a variable $\lambda(t)$, and later finds it to be a constant after all. See Routh E J 1905 *Advanced Dynamics of a System of Rigid Bodies* 6 ed (London: Macmillan), reprinted by Dover, 1955, p.306, and Pars [110], p 249.